

ARITHMETIC AND GEOMETRY OF THE HECKE GROUPS

CHENG LIEN LANG AND MONG LUNG LANG

ABSTRACT. We study the arithmetic and geometry properties of the Hecke group G_q . In particular, we prove that G_q has a subgroup X of index d , genus g with v_∞ cusps, and τ_2 (resp. v_{r_i}) conjugacy classes of elements that are conjugates of S (resp. R^{q/r_i}) if and only if (i) $2g - 2 + \tau_2/2 + \sum_{i=1}^k v_{r_i}(1 - 1/r_i) + v_\infty = d(1/2 - 1/q)$, and (ii) $m_0 = 4g - 4 + \tau_2 + 2v_\infty + \sum_{i=1}^k v_{r_i}(2 - q/r_i) \geq 0$ is a multiple of $q - 2$, (iii) $m \geq 0$. In the case q is odd, (ii) is a consequence of (i).

1. INTRODUCTION

1.1. The (inhomogeneous) Hecke group G_q is defined to be the maximal discrete subgroup of $PSL(2, \mathbb{R})$ generated by S and T , where $\lambda_q = 2\cos(\pi/q)$,

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & \lambda_q \\ 0 & 1 \end{pmatrix}. \quad (1.1)$$

Let $R = ST^{-1}$. Then R has order q and $\{S, R\}$ is a set of independent generators of G_q . Equivalently, G_q is a free product of $\langle S \rangle$ and $\langle R \rangle$. The main purpose of this article is to study the geometric and arithmetic properties of subgroups of finite index of G_q .

1.2. The set of cusps of G_q is $\mathbb{Q}[\lambda_q] \cup \{\infty\}$ if and only if $q = 3, 5$. We will give an inductive procedure (induction on the depth of q -gons) that enables us to generate the set of cusps of G_q (Lemma 3.2). As the index of G_q in $PSL(2, \mathbb{Z}[\lambda_q])$ is infinite, it is important to characterise members of G_q . A simple algorithm that determines whether a matrix of $PSL(2, \mathbb{Z}[\lambda_q])$ belongs to G_q can be found in Proposition 3.7. The algorithm can be implemented in a computer.

1.3. A set of generators $\{x_i\}$ of X is called a set of *independent generators* if X is a free product of the cyclic groups $\langle x_i \rangle$. G_q is a free product of $\langle S \rangle$ and $\langle R \rangle$. By Kurosh's Theorem, every subgroup X of finite index of G_q has a set of independent generators. Proposition 4.4 and Theorem 5.2 demonstrate how arithmetic and geometric properties can be combined to give an inductive procedure for finding a special polygon (fundamental domain) M_X and a set of independent generators I_X for X (the case q is a prime has been done in [LLT1]). In particular, this is applied to the principal congruence subgroup of level 2, the commutator subgroup G'_q and subgroups of index 2 (subsection 5.4).

1.4. As a special case of the Hurwitz-Nielsen realisation problem, Millington [Mi] showed that as long as $d = 3\tau_2 + 4v_3 + 12q + 6t - 12$, then the modular group G_3 possesses a subgroup X of index d , such that $X \setminus \mathbb{H}$ has τ_2 (resp. v_3) elliptic points of order 2 (resp. 3), t cusps, and genus g . We are able to generalise this result to G_q by studying the Hecke-Farey symbols (see Section 6). As mentioned in [K1], the problem of recognising X as a normal subgroup of certain geometric invariants has been left aside in the literature. Our study of this topic starts with some elementary observation of the permutation representations of S and R on the set of cosets G_q/X which we will elaborate more in subsection 1.5.

2010 *Mathematics Subject Classification.* 11F06, 20H10.

Key words and phrases. Hecke groups, congruence subgroups, Kurosh's Theorem, Hurwitz-Nielsen realisation problem, maps and map subgroups.

1.5. We prove that the action of S and R on G_q/X is isomorphic to their action on M_X (see Lemma 7.1) and that the permutation representations of S and R on M_X can be obtained by a simple reading of the special polygon M_X ((7.4) and (7.9)). These two representations $f(S)$ and $f(R)$ carry some important information about X . In particular,

- (i) $f(S)$ and $f(R)$ can be used to determine whether X is normal in G_q and its normaliser in $PSL(2, \mathbb{R})$ (Proposition 8.1 and Discussion 8.2),
- (ii) in the case $q = 3$, $f(S)$ and $f(R)$ can be used to determine whether X is congruence (see Section 11 and [H]),
- (iii) $f(S)$ and $f(R)$ can be used to study the geometric invariants of X (Section 9) and *Dessins d'enfants* (see pp.12 of [HR]). Propositions 9.2-9.4 study the possible realisation of a group Y as a normal subgroup of G_q .

1.6. It is well known that there is a correspondence between the set of *maps* and the set of subgroups of finite index of G_q and that the maps are uniquely determined by their *map subgroups* (see [JS], [CS], [IS]). Let X be a subgroup of G_q of finite index. We give a detailed construction of the map $M(X)$ whose map subgroups are conjugates of X . Both $M(X)$ and X are explicitly given. $\text{Aut } M(X)$ can be determined as well (see Section 10).

2. TESSELLATION OF THE UPPER HALF PLANE

Let D^* denote the $(2, q, \infty)$ triangle with vertices i , $e^{\pi i/q}$ and ∞ . D^* is a fundamental domain of the Coxeter group G_q^* generated by reflections along the sides of D^* . Hecke group G_q is the subgroup of index 2 consists of all the orientation preserving isometries.

Let $\mathbb{H}$ be the union of the upper half plane and the set $\{g(\infty) : g \in G_q^*\}$. The G_q^* translates of D^* form a tessellation $\mathcal{I}^*$ of $\mathbb{H}$ (endowed with the hyperbolic metric) by $(2, q, \infty)$ triangles. The G_q^* translates of i , $e^{\pi i/q}$ and ∞ are called *even vertices*, *odd vertices* and *cusps* (*free vertices*) of $\mathcal{I}^*$ respectively. The G_q^* translates of the hyperbolic line joining i to ∞ (resp. $e^{\pi i/q}$ to ∞) are called *even edges* (resp. *odd edges*) of $\mathcal{I}^*$. The G_q^* translates of the hyperbolic line joining i to $e^{\pi i/q}$ are called *f-edges* of $\mathcal{I}^*$. The hyperbolic line $(0, \infty)$ consists of two even edges. The G_q^* translates of $(0, \infty)$ are called the *even lines* of $\mathcal{I}^*$. The hyperbolic line joining x and y is denoted by (x, y) .

The set of even lines give a tessellation of $\mathbb{H}$ into *ideal q -gons*, that is, hyperbolic q -gons with q cusps. Note that their vertex angle is 0. Each q -gon contains a unique odd vertex.

The *f-edges* form a q -regular tree, where the odd vertices are considered as the set of vertices of this tree. We introduce a vertex of valence two to this q -regular tree at i , denoted by v_0 . Each q -gon P contains a unique vertex v_P of this tree. The *depth* of P denoted by $d(P)$ is defined to be the distance between v_0 and v_P (the distance between adjacent vertices is 1).

3. CUSPS AND REDUCED FORMS

The main purpose of this section is to determine whether a matrix of $PSL(2, \mathbb{Z}[\lambda_q])$ belongs to G_q (Proposition 3.7). Lemma 3.2 gives the set of cusps of G_q .

3.1. Reduced forms of cusps of G_q . The set of cusps of G_q is a subset of $\mathbb{Q}(\lambda_q) \cup \{\infty\}$. Let x be a cusp. We say $x = a/b$ is in *reduced form* if

- (i) there exists c/d such that $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \in G_q$, and (ii) $b \geq 0$.

Let g be given as in (i) of the above. Since $w = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ normalises G_q , $-wgw^{-1} \in G_q$. Hence a/b is in reduced form if and only if $-a/b$ is in reduced form. In the case $a, b \geq 0$, it is also easy to see that a/b is in reduced form if and only if $-b/a$ is in reduced form (study $S^{-1}g$) if and only if b/a is in reduced form (study $w(Sg)w^{-1}$).

3.2. Construction of reduced forms. We give some basics about reduced forms.

Lemma 3.1. *Suppose that $w \neq \infty$. The reduced form of w is unique. The reduced form of ∞ is either $-1/0$ or $1/0$.*

Proof. Let a/b and a'/b' be the reduced forms of w . It is clear that $ab' = a'b$. Following our definition of reduced form, G_q contains the following two elements, $\begin{pmatrix} a & x \\ b & y \end{pmatrix}$ and $\begin{pmatrix} a' & u \\ b' & v \end{pmatrix}$. An easy calculation gives $\begin{pmatrix} a & x \\ b & y \end{pmatrix}^{-1} \begin{pmatrix} a' & u \\ b' & v \end{pmatrix} = \begin{pmatrix} a'y - b'x & yu - xv \\ 0 & av - bu \end{pmatrix} \in G_q$. This element fixes ∞ . Since the stabiliser of ∞ in G_q is generated by T (see (1.1)), we have the following.

$$\begin{pmatrix} a & x \\ b & y \end{pmatrix}^{-1} \begin{pmatrix} a' & u \\ b' & v \end{pmatrix} = \begin{pmatrix} 1 & m\lambda_q \\ 0 & 1 \end{pmatrix}. \quad (3.1)$$

It follows from (3.1) that $a = a'$ and that $b = b'$. This completes the proof of the lemma. $\square$

Lemma 3.2. *Let P be an ideal q -gon with cusps $\{c_1, c_2, \dots, c_q\}$ arranged in increasing order (if ∞ is a cusp, then $c_1 = -\infty = -1/0$ if P lies in the left half plane and $c_q = \infty = 1/0$ if P lies in the right half plane). Let a_i/b_i be the reduced form of c_i and $\frac{a_0}{b_0} = \frac{-a_q}{-b_q}$. Then*

$$a_i = \lambda_q a_{i-1} - a_{i-2} \text{ and } b_i = \lambda_q b_{i-1} - b_{i-2} \text{ for } 2 \leq i \leq q-1. \quad (3.2)$$

Proof. Without loss of generality, we assume that P lies in the right half plane. Let $A = T^{-1}S = \begin{pmatrix} \lambda_q & 1 \\ -1 & 0 \end{pmatrix} \in G_q$ and $w_i = \begin{pmatrix} a_{i-1} & a_{i-2} \\ b_{i-1} & b_{i-2} \end{pmatrix}$, $2 \leq i \leq q$. Note that $Ae^{(q-1)\pi i/q} = e^{(q-1)\pi i/q}$ and that A is a counter-clockwise rotation about $e^{(q-1)\pi i/q}$ of angle $2\pi/q$.

(A) Suppose that the depth of P is 1. Then (c_1, c_q) is $(0, \infty)$ and $e^{\pi i/q}$ is the odd vertex of P . We apply mathematical induction as follows.

- (i) It is clear that $w_2 = S^{-1} \in G_q$, $w_2 A w_2^{-1}$ fixes $e^{\pi i/q}$ and that $w_3 = w_2 A \in G_q$.
- (ii) Suppose that $w_{i-1} \in G_q$, $w_{i-1} A w_{i-1}^{-1}$ fixes $e^{\pi i/q}$, and that $w_i = w_{i-1} A \in G_q$.

By (ii), $w_i A w_i^{-1} = w_{i-1} A w_{i-1}^{-1}$ fixes $e^{\pi i/q}$ and $w_i A w_i^{-1}$ is a counter-clockwise rotation about $e^{\pi i/q}$ of angle $2\pi/q$. Hence $w_i A w_i^{-1}$ sends $(c_{i-1}, c_{i-2}) = (a_{i-1}/b_{i-1}, a_{i-2}/b_{i-2})$ to $(c_i, c_{i-1}) = (a_i/b_i, a_{i-1}/b_{i-1})$ and c_i is given by the first column of the following matrix

$$w_i A w_i^{-1} w_i = \begin{pmatrix} a_{i-1} & a_{i-2} \\ b_{i-1} & b_{i-2} \end{pmatrix} A = \begin{pmatrix} \lambda_q a_{i-1} - a_{i-2} & a_{i-1} \\ \lambda_q b_{i-1} - b_{i-2} & b_{i-1} \end{pmatrix} \in G_q. \quad (3.3)$$

To be more precise, $(\lambda_q a_{i-1} - a_{i-2})/(\lambda_q b_{i-1} - b_{i-2}) = a_i/b_i = c_i$. Since $c_i > c_{i-1}$, $\det w_i A > 0$, and $a_k, b_k \geq 0$ for all k , one has $\lambda_q b_{i-1} - b_{i-2} > 0$. Hence $(\lambda_q a_{i-1} - a_{i-2})/(\lambda_q b_{i-1} - b_{i-2})$ is in reduced form (subsection 3.1). Since the reduced form of c_i is unique, one has $a_i = \lambda_q a_{i-1} - a_{i-2}$, $b_i = \lambda_q b_{i-1} - b_{i-2}$ and $w_{i+1} = w_i A \in G_q$. By induction, we conclude that if the depth of P is one, then $a_i = \lambda_q a_{i-1} - a_{i-2}$ and $b_i = \lambda_q b_{i-1} - b_{i-2}$.

(B) Suppose that the depth of P is 2 or more. It follows that (c_1, c_q) is a side of a q -gon of depth $d(P) - 1$. By induction on the depth of P , we may assume that the reduced forms of $c_1 = a_1/b_1$ and $c_q = a_q/b_q$ are known and that $w_2 = \begin{pmatrix} a_1 & -a_q \\ b_1 & -b_q \end{pmatrix} \in G_q$. Note that $v_0 = w_2 e^{(q-1)\pi i/q}$ is the odd vertex of P and that $w_2 A w_2^{-1}$ fixes v_0 . We may now apply induction as above (see case (A)) and conclude that $a_i = \lambda_q a_{i-1} - a_{i-2}$ and that $b_i = \lambda_q b_{i-1} - b_{i-2}$. This completes the proof of the lemma. $\square$

Lemma 3.2 tells us how to generate the reduced forms of all positive cusps starting with $(0/1, 1/0)$ and all negative cusps starting with $(-1/0, 0/1)$. Namely,

$$\frac{a_i}{b_i} = \frac{\lambda_q a_{i-1} - a_{i-2}}{\lambda_q b_{i-1} - b_{i-2}}, \quad i = 2, 3, \dots, q-1. \quad (3.4)$$

Equation (3.4) allows us to write the c_i 's ($1 \leq i \leq q$) in terms of c_1 and c_q which generalises the construction of the Farey sequence. In the case $q = 6$, $\lambda_6 = \sqrt{3}$, the reduced forms of

the c_i 's ($1 \leq i \leq 6$) in terms of $c_1 = a_1/b_1$ and $c_6 = a_6/b_6$ are given by

$$\frac{a_1}{b_1}, \frac{\sqrt{3}a_1 + a_6}{\sqrt{3}b_1 + b_6}, \frac{2a_1 + \sqrt{3}a_6}{2b_1 + \sqrt{3}b_6}, \frac{\sqrt{3}a_1 + 2a_6}{\sqrt{3}b_1 + 2b_6}, \frac{a_1 + \sqrt{3}a_6}{b_1 + \sqrt{3}b_6}, \frac{a_6}{b_6}. \quad (3.5)$$

The continuous solution of the equation $b_i = \lambda_q b_{i-1} + b_{i-2}$ ($b_i \geq 0$, $2 \leq i \leq q-1$) is

$$f(x) = \frac{b_1 + b_q \cos(\pi/q)}{\sin(\pi/q)} \sin(\pi x/q) - b_q \cos(\pi x/q) = A \sin(\pi x/q + w), \quad f(i) = b_i, \quad (3.6)$$

for some A and w . Note that $f(x)$ is a periodic function of period $2q$, $f(0) = -b_q < 0$, and $f(1) = b_1 > 0$. Further, $f(x)$ is concave down in the interval $[0, q]$. As a consequence,

$$b_i \geq b_1 \text{ and } b_i \geq b_q \text{ for all } i. \quad (3.7)$$

Lemma 3.3. *Let a/b be in reduced form. Suppose that $ab \neq 0$. Then $|a| \geq 1$ and $b \geq 1$.*

Proof. Since a/b is in reduced form, $a/b = x_i$ is a cusp of some q -gon P . Let $\{x_1, x_2, \dots, x_q\}$ be the set of cusps of P . We prove that $b \geq 1$ by induction on the depth of P . In the case the depth of P is 1, the set of cusps of P is either $\{0/1, 1/\lambda_q, \dots, 0/1\}$ if P lies in the right half plane or $\{-0/1, \dots, -1/\lambda_q, 0/1\}$ if P lies in the left half plane. By (3.7), $b \geq 1$. Suppose that our assertion holds when the depth of P is n . We now consider the case $d(P) = n+1$ and that a/b is a cusp of P . In the case that a/b is either x_1 or x_q , a/b is also a cusp of a q -gon of depth n . By inductive hypothesis, $b \geq 1$. Hence we shall assume that $a/b = x_i$, where $2 \leq i \leq q-1$. By (3.7), b is larger than the denominators of the reduced forms of x_1 and x_q . As x_1 and x_q are cusps of a q -gon of depth n , their denominators are at least 1 by inductive hypothesis. Hence $b \geq 1$. Since a/b is in reduced form, $b/|a|$ is also in reduced form (see subsection 3.1). Repeat the above argument, one has $|a| \geq 1$. $\square$

3.3. Pseudo Euclidean algorithm. Let $x, y \in \mathbb{Z}[\lambda_q]$. Suppose that $y \neq 0$. There exists a unique integer $m \in \mathbb{Z}$ such that $x = y(m\lambda_q) + r$, $-|y\lambda_q|/2 < r \leq |y\lambda_q|/2$. We call such an algorithm *pseudo Euclidean* (PEA). Let $a, b \in \mathbb{Z}[\lambda_q]$, where $ab \neq 0$. Apply the pseudo Euclidean algorithm repeatedly,

$$\begin{aligned} a &= b(m_0\lambda_q) + r_1, \\ b &= r_1(m_1\lambda_q) + r_2, \\ &\vdots \\ r_{k-1} &= r_k(m_k\lambda_q) + r_{k+1}. \end{aligned} \quad (3.8)$$

Let $r_0 = b$. If the (PEA) terminates, that is $r_n \neq 0$, $r_{n+1} = 0$ for some n , we define

$$(a, b)_q = |r_n|. \quad (3.9)$$

If the (PEA) does not terminate, we define $(a, b)_q = 0$. Define further that $(a, 0)_q = (0, a)_q = |a|$. One sees easily that (i) $(a, b)_q = (b, a)_q$, (ii) $(a, b)_q = (-a, -b)_q = (-a, b)_q = (a, -b)_q$.

Lemma 3.4. *Let a/b be in reduced form. If $b = 1$, then $a = m\lambda_q$ for some $m \in \mathbb{Z}$.*

Proof. Suppose that $a \neq 0$. Since a/b is in reduced form, the (PEA) (see (3.8)) implies that r_1/b is in reduced form and that $|r_1| < b\lambda_q/2 < b$. Since $b = 1$, one has $|r_1| < 1$. By Lemma 3.3, $r_1 = 0$. Hence $a = m\lambda_q$ for some $m \in \mathbb{Z}$. This completes the proof of the lemma. $\square$

Remark. The converse of Lemma 3.4 is not true as G_4 and G_6 possess infinitely many reduced forms $m\lambda_q/b$, where $m, b \in \mathbb{N}$, $b > 1$.

Lemma 3.5. *Let c/d and a/b be the reduced forms of x and y respectively, where $x < y$. Then $g = \begin{pmatrix} a & c \\ b & d \end{pmatrix} \in G_q$ if and only if (x, y) is an even line if and only if $ad - bc = 1$.*

Proof. (i) Suppose that $g \in G_q$. Then $g(0, \infty)$ is an even line. Equivalently, (x, y) is an even line. (ii) Suppose that (x, y) is an even line. Then $A(0, \infty) = (x, y)$ for some $A \in G_q$. An easy study of Lemma 3.1 and the matrix form of $A(0, \infty) = (x, y)$ shows that $ad - bc = 1$. (iii) Suppose that $ad - bc = 1$. Since a/b and c/d are in reduced forms, G_q contains elements

of the following forms, $\begin{pmatrix} a & x \\ b & y \end{pmatrix}$ and $\begin{pmatrix} c & u \\ d & v \end{pmatrix}$. An easy calculation show that $\begin{pmatrix} a & x \\ b & y \end{pmatrix}^{-1} \begin{pmatrix} c & u \\ d & v \end{pmatrix} = \begin{pmatrix} cy - dx & * \\ 1 & * \end{pmatrix}$. By Lemma 3.4, one has $cy - dx = m\lambda_q$ for some $m \in \mathbb{Z}$. It follows that $\begin{pmatrix} a & x \\ b & y \end{pmatrix}^{-1} g = \begin{pmatrix} a & x \\ b & y \end{pmatrix}^{-1} \begin{pmatrix} a & c \\ b & d \end{pmatrix} = \begin{pmatrix} 1 & m\lambda_q \\ 0 & 1 \end{pmatrix} \in G_q$. Hence $g \in G_q$. $\square$

Lemma 3.6. *Suppose that $(a, b)_q = r \neq 0$. Then there exists some $g \in G_q$ such that $g \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} r \\ 0 \end{pmatrix}$. In particular, $(a/r, b/r)_q = 1$ and $\frac{a/r}{|b/r|}$ is in reduced form.*

Proof. The lemma follows from (3.8), (3.9) and the observation that the matrix form of the equation $x = y(m\lambda_q) + z$ is $\begin{pmatrix} 1 & -m\lambda_q \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} z \\ y \end{pmatrix}$, where $\begin{pmatrix} 1 & -m\lambda_q \\ 0 & 1 \end{pmatrix} \in G_q$. $\square$

Proposition 3.7. *Let $A = \begin{pmatrix} a & c \\ b & d \end{pmatrix} \in PSL(2, \mathbb{Z}[\lambda_q])$. Then $A \in G_q$ if and only if $(a, b)_q = (c, d)_q = 1$. In particular, if $(a, b)_q < 1$ or $|b| < 1$, then A is not in G_q .*

Proof. Suppose that $A \in G_q$. Then $a/|b|$ is in reduced form (see subsection 3.1). By (3.8), $r_n/|r_{n+1}|$ is in reduced form for every n . By Lemma 3.3, $|r_n| \geq 1$ and $|r_{n+1}| \geq 1$ whenever $r_n r_{n+1} \neq 0$. Since a and b are finite and $\lambda_q/2 < 1$, an easy observation of (3.8) implies that there exists some m such that $|r_k| < 1$ whenever $k \geq m$. Hence there exists a d such that $r_{d+1} = 0$ and that $r_d \neq 0$. By Lemma 3.1, $r_d = \pm 1$. Equivalently, $(a, b)_q = 1$. Since $A \in G_q$, $A^{-1}S \in G_q$. Since the transpose of S and T are members of G_q , the transpose of $A^{-1}S$ is also an element of G_q , the first column of the transpose of $A^{-1}S$ is $\begin{pmatrix} c \\ d \end{pmatrix}$. This implies that $c/|d|$ is in reduced form. Similar to the above, one can show that $(c, d)_q = 1$.

Conversely, suppose that $(a, b)_q = (c, d)_q = 1$. Replace A by $-A$ if necessary, we may assume that $d \geq 0$. Replace A by wAw^{-1} if necessary (see subsection 3.1 for w), we may assume that $b \geq 0$ and that $d \geq 0$. By Lemma 3.6, both a/b and c/d are in reduced forms. By Lemma 3.5, we have $A \in G_q$. This completes the proof of the lemma. $\square$

Example 3.8. Let $q = 5$ and let $\lambda = \lambda_5$. Note that $\lambda^2 = \lambda + 1$. $\begin{pmatrix} 4\lambda - 1 & \lambda + 1 \\ 3 & \lambda \end{pmatrix}$ is not an element of G_5 as $(4\lambda - 1, 3)_5 = \lambda - 1 < 1$.

4. HECKE-FAREY SYMBOLS AND SPECIAL POLYGONS

4.1. r -clusters. Let Φ be the hyperbolic triangle with vertices $0, e^{\pi i/q}$ and ∞ . Φ is a fundamental domain of G_q . The G_q translates of Φ are called *special triangles*. For each divisor r of q ($1 \leq r < q$), set $\Phi_r = \Phi \cup R\Phi \cup R^2\Phi \cup \dots \cup R^{r-1}\Phi$. Φ_r is a union of r copies of special triangles. These r special triangles meet at the odd vertex $e^{\pi i/q}$. The G_q translates of Φ_r are called the *r -clusters* (a 1-cluster is a special triangle). Let Δ_r be an r -cluster. It is clear that (i) Δ_r has $(r + 1)$ cusps and one odd vertex y , (ii) the boundary of Δ_r has r even lines and two odd edges, (iii) the two odd edges of Δ_r meet each other at y with vertex angle $2r\pi/q$. The cusps are called the *free vertices* of Δ_r .

Let $\{v_1 = 0/1, v_2, \dots, v_{q-1}, v_q = 1/0\}$ be the set of cusps of the depth 1 q -gon in the right half plane where the v_i 's are arranged in increasing order. Following the definition of Φ_r , the set of free vertices (cusps) of Φ_r is

$$\Phi_r : \{0 = v_1, v_2, \dots, v_r, v_q = \infty\}. \quad (4.1a)$$

The odd vertex of Φ_r is $e^{\pi i/q}$. The odd edges are $(e^{\pi i/q}, v_r)$ and $(e^{\pi i/q}, v_q) = (e^{\pi i/q}, \infty)$. Note that Φ_r is convex as it is the convex hull of $\{v_1, v_2, \dots, v_r, e^{\pi i/q}, v_q\}$. Let Ψ_r be the mirror image of Φ_r (with respect to the y -axis). The set of free vertices of Ψ is

$$\Psi_r : \{-\infty = -v_q, -v_r, \dots, -v_2, -v_1 = 0\}. \quad (4.1b)$$

The odd vertex of Ψ_r is $y = e^{(q-1)\pi i/q}$ and the odd edges of Ψ_r are $(y, -v_r)$ and $(y, -v_q)$. Let A be given as in Lemma 3.2. It is clear that $\Psi_r = A^{1-r}S\Phi$.

Example 4.1. Let $q = 6$, $\lambda = \lambda_6 = \sqrt{3}$. See **Figure 1a** for Φ_2 and **Figure 1b** for Ψ_2 .




Figure 1a




Figure 1b

4.2. Special polygons. A convex hyperbolic polygon P of $\mathbb{H}$ is a union of some q -gons and a finite number of r_i -clusters ($r_i|q$, $1 \leq r_i < q$). The q -gons and the r_i -clusters of P are called the *tiles*. The tiles intersect each other (if any) at either free vertices (cusps) or even lines. A *special polygon* $M_X = (P, I_X)$ of $\mathbb{H}$ is a convex hyperbolic polygon P together with a set of side pairings I_X satisfying the rules below.

- (S1) An odd edge e is always paired with an odd edge f (in the same r -cluster) and makes an internal angle $2r\pi/q$ with f . The vertex where e and f meet is an *odd vertex* of P . Both e and f are considered as sides of P , and are called its *odd sides*.
- (S2) Let e and f be two even edges in the boundary of P forming an even line. Then either (i) e is paired with f , both e and f are considered as sides of P , and are called its *even sides*, the point where e and f meet is an *even vertex* of P , or (ii) e and f form a *free side* of P , and this free side is paired with another free side of P .
- (S3) 0 and ∞ are vertices of P .

Let $M_X = (P, I_X)$ be a special polygon. The cusps in P are called the *free vertices* of M_X .

4.3. Hecke-Farey sequences and symbols. A *Hecke-Farey sequence* is a finite sequence of cyclically arranged numbers in increasing order $\{-\infty, x_0, x_1, \dots, x_n, \infty\}$ such that

- (a) $x_i \in \mathbb{Q}[\lambda_q] = \mathbb{Q}(\lambda_q)$, $x_i = 0$ for some i , $0 \leq i \leq n$,
- (b) $x_i = a_i/b_i$ is in reduced form for every i ($x_{-1} = -\infty = -1/0$ and $x_{n+1} = \infty = 1/0$),
- (c) if $a_{i+1}b_i - a_ib_{i+1} \neq 1$, then there exists an element $g \in G_q$ and an r -cluster Φ_r ($r|q$, $1 < r < q$) such that $g(v_r) = x_i$, $g(v_q) = x_{i+1}$ (see (4.1a) for v_r and v_q).
- (d) $\{x_i, x_{i+1}\}$ is called an *interval*. Intervals described as in (c) are called *r -intervals*. $\{x_i, x_{i+1}\}$ is called an *ordinary interval* if $a_{i+1}b_i - a_ib_{i+1} = 1$. By Lemma 3.5, if $\{x_i, x_{i+1}\}$ is an ordinary interval, then the hyperbolic line (x_i, x_{i+1}) is an even line.

One can show easily that that the g 's and r 's in (c) of the above are unique (the only element of G_q that fixes two or more points of $\mathbb{H}$ is the identity). A *Hecke-Farey symbol* is a Hecke-Farey sequence together with an additional *side pairing* on each consecutive pair of x_i 's. To avoid triviality, we insist that a Hecke-Farey symbol must have at least two distinct side pairings (see Discussion 4.5). In the case $a_{i+1}b_i - a_ib_{i+1} \neq 1$, the additional side pairing of the r -interval $\{x_i, x_{i+1}\}$ is denoted by

$$x_i \underset{e_{r(g)}}{\smile} x_{i+1}, \quad (4.2)$$

where g and r are given as in (c) of the above. In the case $a_{i+1}b_i - a_ib_{i+1} = 1$, the additional side pairing of the interval $\{x_i, x_{i+1}\}$ can be either one of the following three types :

$$x_i \underset{\circ}{\smile} x_{i+1}, \quad x_i \underset{\bullet}{\smile} x_{i+1}, \quad x_i \underset{a}{\smile} x_{i+1}, \quad (4.3)$$

where a is a natural number. Each natural number a occurs exactly twice or not at all. The actual values of the a 's are unimportant, it is the pairing induced on the consecutive pairs that matters. We shall now give a detailed description of these side pairings.

- (i) Let $\{x_i, x_{i+1}\}$ be an r -interval and let g, Φ_r be given as in (c) of the above. By (c), one has $g(v_r) = x_i$ and $g(v_q) = x_{i+1}$. Set $g(e^{\pi i/q}) = y$.

$$x_i \underset{e_r(g)}{\smile} x_{i+1} \quad (4.4)$$

is the side pairing that fixes y and sends the odd edge (y, x_{i+1}) to (y, x_i) . Since R is a counterclockwise rotation about $e^{\pi i/q}$ of angle $2\pi/q$ and (y, x_i) and (y, x_{i+1}) form an internal angle $2r\pi/q$, this side pairing must be $gR^r g^{-1} \in G_q$ (the only element of G_q that fixes two or more points of $\mathbb{H}$ is the identity).

- (ii) Let $\{x_i, x_{i+1}\}$ be an ordinary interval. By (d) of the above, (x_i, x_{i+1}) is an even line. Hence there exists a unique $A \in G_q$ such that $A(0) = x_i$ and $A(\infty) = x_{i+1}$. Set $A(e^{\pi i/q}) = y$. (y, x_{i+1}) and (y, x_i) are odd edges.

$$x_i \underset{\bullet}{\smile} x_{i+1} \quad (4.5)$$

is the side pairing that fixes y and sends the odd edge (y, x_{i+1}) to (y, x_i) . It is clear that this side pairing must be

$$h = ARA^{-1} = \begin{pmatrix} a_{i+1} & a_i \\ b_{i+1} & b_i \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & \lambda_q \end{pmatrix} \begin{pmatrix} a_{i+1} & a_i \\ b_{i+1} & b_i \end{pmatrix}^{-1} \in G_q. \quad (4.5a)$$

- (iii) Let $\{x_i, x_{i+1}\}$ be an ordinary interval. By (d) of the above, (x_i, x_{i+1}) is an even line. Hence there exists a unique $A \in G_q$ such that $A(0) = x_i$ and $A(\infty) = x_{i+1}$. Set $A(\sqrt{-1}) = y$. (y, x_{i+1}) and (y, x_i) are even edges.

$$x_i \underset{\circ}{\smile} x_{i+1} \quad (4.6)$$

is the side pairing that fixes y and sends the even edge (y, x_{i+1}) to (y, x_i) . It is clear that this side pairing must be

$$w = ASA^{-1} = \begin{pmatrix} a_{i+1} & a_i \\ b_{i+1} & b_i \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} a_{i+1} & a_i \\ b_{i+1} & b_i \end{pmatrix}^{-1} \in G_q. \quad (4.6a)$$

- (iv) Let $\{x_i, x_{i+1}\}$ and $\{x_j, x_{j+1}\}$ be two ordinary intervals. By (d) of the above, (x_j, x_{j+1}) and (x_i, x_{i+1}) are even lines.

$$x_i \underset{\alpha}{\smile} x_{i+1}, \quad x_j \underset{\alpha}{\smile} x_{j+1} \quad (4.7)$$

is the side pairing that sends x_{j+1} to x_i and x_j to x_{i+1} (equivalently, pairs the even lines (x_j, x_{j+1}) and (x_i, x_{i+1})). It is clear that this side pairing must be

$$k = \begin{pmatrix} a_j & -a_{j+1} \\ b_j & -b_{j+1} \end{pmatrix} \begin{pmatrix} a_{i+1} & a_i \\ b_{i+1} & b_i \end{pmatrix}^{-1} \in G_q. \quad (4.7a)$$

Definition 4.2. The intervals in (ii) are called *odd intervals*, the intervals in (iii) are called *even intervals*, and the intervals in (iv) are called *free intervals*.

Example 4.3. Let $q = 6$, $\lambda = \lambda_6 = \sqrt{3}$, $g = \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix}$. The following is a Hecke-Farey symbol (see Figure 2).

$$M_X = \{-\infty \underset{\circ}{\smile} 0/1 \underset{\circ}{\smile} 1/2\lambda \underset{e_2(g)}{\smile} 1/\lambda \underset{1}{\smile} \lambda/2 \underset{1}{\smile} 2/\lambda \underset{1}{\smile} \lambda/1 \underset{\bullet}{\smile} \infty\}. \quad (4.8)$$

Proof. The set of cusps of Φ_2 is $\{0/1, 1/\lambda, \infty\}$. One must verify that the interval $\{1/2\lambda, 1/\lambda\}$ satisfies (c) of the above. This can be checked easily as $g(1/\lambda) = 1/2\lambda$ and $g(\infty) = 1/\lambda$. Note that $g\Phi_2$ is the 2-cluster with odd vertex y . $\square$




Figure 2

4.4. Special polygons and Hecke-Farey symbols. We now relate Hecke-Farey symbols to special polygons. It is a generalisation of a theorem of [K3] and is proved in an analogous way (see Example 4.3 and Figure 2 for an example).

Proposition 4.4. *There is a one to one correspondence between the set of special polygons and the set of Hecke-Farey symbols.*

Proof. Let $M_X = (P, I_X)$ be a special polygon. The free vertices (cusps) of P form a sequence $\{-\infty = x_{-1}, x_0, x_1, \dots, x_n, \infty = x_{n+1}\}$. Let a_k/b_k be the reduced form of x_k and let x_i, x_{i+1} be two consecutive terms. Suppose that $a_{i+1}b_i - a_i b_{i+1} \neq 1$. By Lemma 3.5, (x_i, x_{i+1}) is not an even line of P . Following the definition of convex hyperbolic polygons (see (S1)-(S3) of subsection 4.2), one has (i) x_i and x_{i+1} are the end points of two odd edges $e = (y, x_i)$ and $f = (y, x_{i+1})$ of P that meet at an odd vertex y , (ii) e and f are the odd edges of an r -cluster $g\Phi_r$ for some g and r . As a consequence, x_i and x_{i+1} satisfy (c) of subsection 4.3 and gives an r -interval $\{x_i, x_{i+1}\}$. Hence the sequence

$$F = \{-\infty = x_{-1}, x_0, x_1, \dots, x_n, \infty = x_{n+1}\} \quad (4.9)$$

is a Hecke-Farey sequence. In the case $\{x_i, x_{i+1}\}$ is an r -interval, following our notation above and (S1) of subsection 4.2, $e = (y, x_i)$ and $f = (y, x_{i+1})$ are paired by an element of I_X . As this side pairing must be unique, it has to be $gR^r g^{-1}$ (see (4.4)). Hence the r -interval must carry the side pairing given by (4.4). In the case $\{x_i, x_{i+1}\}$ is an ordinary interval of F , similar study of (S1)-(S3) of subsection 4.2 implies that the side pairing in I_X that does the pairing for $\{x_i, x_{i+1}\}$ is just the side pairing given by (4.3). This makes F into a Hecke-Farey symbol.

Conversely, given a Hecke-Farey symbol, one can construct a special polygon with free vertices corresponding to the points of the Hecke-Farey sequence, odd vertices where the symbols $e_r(g)$ and $\bullet$ occur and even vertices where the symbol $\circ$ occurs and side pairings as determined by (i)-(iv) of subsection 4.3. $\square$

Discussion 4.5. Note that $F_1 = \{-\infty \underset{\circ}{\cup} 0/1 \underset{\circ}{\cup} \infty\}$ and $F_2 = \{-\infty \underset{1}{\cup} 0/1 \underset{1}{\cup} \infty\}$ are not Hecke-Farey symbols since a Hecke-Farey symbol must have at least two side pairings (see subsection 4.3) whereas F_1 has only one side pairing S and F_2 has only one side pairing I_2 . It is clear that F_1 and F_2 do not correspond to any special polygon. As a consequence, Proposition 4.4 is no longer true if one does not insist that a Hecke-Farey symbol must carry at least two side pairings.

5. POINCARÉ'S POLYGONAL THEOREM AND INDEPENDENT GENERATORS

Theorem 5.1. *If $M_X = (P, I_X)$ is a special polygon, then the set of side pairings I_X generates independently a group $G \subseteq G_q$ such that P is a fundamental domain of G .*

Proof. The stabilisers of the cusps of P in G are generated by conjugates of T^n for some $n \in \mathbb{N}$. Let $v \in P$ be an elliptic point of P . The stabiliser $G_v = \langle \tau \rangle$ of v in G is generated by either a conjugate of S or a conjugate of R^r for some r , where $r|q$. It is clear from our construction of P that (i) v is the intersection of two edges e_1 and e_2 of P (see (S1) and

(S2) of subsection 4.2), (ii) the sides e_1 and e_2 make an internal angle π or $2r\pi/q$ at v , and (iii) we may assume that τ fixes v and sends e_1 to e_2 . In summary,

- (i) the stabiliser of a cusp of P in G is generated by a parabolic element,
- (ii) e_1 and e_2 make an internal angle π or $2r\pi/q$ at v , where $r|q$ and $r < q$,
- (iii) τ is the side pairing that fixes v and sends e_1 to e_2 .

Hence P is a Poincaré polygon. By Poincaré's polygonal theorem, P is a fundamental domain of G and I_X is a set of independent generators of G (see pp 223 of [Ma]). $\square$

Let X be a subgroup of finite index of G_q . An *admissible fundamental domain* is a special polygon $M_X = (P, I_X)$ such that P is a fundamental domain of X and I_X is a set of independent generators of X .

Theorem 5.2. *Let X be a subgroup of finite index of G_q . Then X has an admissible fundamental domain M_X .*

Proof. The tessellation $\mathcal{I}^*$ (see subsection 2.1) induces a tessellation of the surface $X \setminus \mathbb{H}$. Let v be an odd vertex of a tile T_0 of $X \setminus \mathbb{H}$. The stabiliser $X_v \subseteq X$ is cyclic of order d , where d is a divisor of q . In the case $d > 1$, v is the odd vertex of a q/d -cluster T_0 . Note that the two odd sides of T_0 must be paired by some members of X_v . In the case $d = 1$, v is the odd vertex of a q -gon T_0 . In summary, the tiles of $X \setminus \mathbb{H}$ are q -gons and q/d -clusters, where $d|q$. If $X \setminus \mathbb{H}$ has an elliptic point of order 2 on the boundary of T_0 , then the two even edges incident to the elliptic point are paired by an element of X and forming an edge of $X \setminus \mathbb{H}$. It is clear that these tiles intersect each other (if any) at either cusps or even lines as the two odd sides of a q/d -cluster must be paired by some elements in X_v .

Finding a fundamental domain for X which is a special polygon amounts to cutting the surface $X \setminus \mathbb{H}$ into its tiles (q -gons and q/d -clusters) and develop these tiles on $\mathbb{H}$ so that 0 and ∞ are vertices of the polygon (see (S3) of subsection 4.2).

(A) Let $p : \mathbb{H} \rightarrow X \setminus \mathbb{H}$ be the projection map and let $(0, \infty)$ be the even line joining 0 and ∞ . $p((0, \infty))$ lies on the boundary of some tile T of $X \setminus \mathbb{H}$. We develop T to $\mathbb{H}$ so that $p((0, \infty))$ is developed to $(0, \infty)$. The other tiles of $X \setminus \mathbb{H}$ are then developed onto $\mathbb{H}$ in an inductive manner, piece by piece, where each new piece is adjacent to a tile that is already been developed.

(B) The determination of the first tile P_0 .

- (i) Let $D_1 = \{\Phi_d : \text{the two odd edges of } \Phi_d \text{ are paired by some elements of } X\}$ and $D_2 = \{\Psi_d : \text{the two odd edges of } \Psi_d \text{ are paired by some elements of } X\}$ (see subsection 4.1 for Φ_d and Ψ_d). If $D_1 \neq \emptyset$, let P_0 be the smallest Φ_d (in area) of D_1 . If $D_1 = \emptyset$ and $D_2 \neq \emptyset$, let P_0 be the smallest Ψ_r (in area) of D_2 . If $D_1 \cup D_2 = \emptyset$, let P_0 be the depth one q -gon that lies in the right half plane. Note that 0 and ∞ are vertices of P_0 (see (S3) of subsection 4.2).
- (ii) The reduced forms of the vertices of P_0 can be determined by Lemma 3.2. Note that the two odd edges of P_0 (if any) are paired by some elements of X .

(C) Let e be a side (even line) of P_0 .

- (i) If there exists another side f of P_0 such that the element $g \in G_q$ which pairs e and f is in X , then we call e and f *paired sides* and add g to the generating set I_X of X . The side pairing g can be determined by (4.7).
- (ii) If the element g in G_q that pairs the two even edges of e is in X , then we call e a *paired side* and g is put into the generating set I_X of X . Note that g has order 2. Such g can be determined by (4.6).
- (iii) Let D be the collection of all d -clusters ($d < q$) attached to e and let $D_e = \{T \in D : \text{the two odd edges of } T \text{ are paired by some elements of } X\}$. If D_e is not empty, then we attach T_0 to P_0 (along e) to form a new polygon $P_1 = P_0 \cup T_0$, where T_0 is the smallest d -cluster (in area) of D_e . The two odd edges of T_0 paired by $g \in X$




Figure 3a




Figure 3b

are called *paired sides* and g is put into the generating set I_X of X as before. Note that g can be determined by (4.4) and (4.5).

- (iv) If e does not satisfy any of the above, e is called an *unpaired side* of P_0 .
- (v) It is clear that only one of the four cases can occur to e .

(D) The inductive step.

We have a polygon P_n with some paired sides and some unpaired sides. If there is no unpaired sides, we are done and P_n with the side pairings is a special polygon of X . Otherwise choose any one of the unpaired sides e and adjoin a q -gon T_n along e to form a new polygon $P_{n+1} = P_n \cup T_n$. The reduced forms of the vertices of T_n can be obtained by applying Lemma 3.2. P_{n+1} has $q - 1$ new sides (coming from T_n). We check for cases (i), (ii) and (iii) of (C) for each of the new sides. If after checking (i)-(iii), P_{n+1} has no more unpaired sides, we are done. Otherwise we continue inductively.

Since the index of X is finite, this process must stop after a finite number of steps. The polygon P together with the set of side pairings I_X is a special polygon. By Theorem 5.1, (P, I_X) is an admissible fundamental domain of X . $\square$

5.1. Two subgroups of the Hecke group G_6 of index 3. (i) Let $\lambda = \lambda_6 = \sqrt{3}$ and let M_X be a special polygon of X given as follows (see Figure 3a).

$$M_X = \{-\infty \underset{\circ}{\smile} 0 \underset{\circ}{\smile} 1/\lambda \underset{\circ}{\smile} \lambda/2 \underset{e_3(1)}{\smile} \infty\}. \quad (5.1)$$

To determine the side pairing $e_3(1)$, we follow the notation of (i) subsection 4.3. One has $r = 3$, $g = 1$, $x_i = \lambda/2$ and that $x_{i+1} = 1/0$. By (4.4), one has $e_3(1) = R^3$. X is a normal subgroup of index 3 (see Discussion 8.2 for normality). The remaining side pairings can be determined easily by (4.6). In summary, a set of independent generators is given by

$$X = \left\langle S, \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix} S \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix}^{-1}, \begin{pmatrix} \lambda & 1 \\ 2 & \lambda \end{pmatrix} S \begin{pmatrix} \lambda & 1 \\ 2 & \lambda \end{pmatrix}^{-1}, \begin{pmatrix} 0 & 1 \\ -1 & \lambda \end{pmatrix}^3 \right\rangle. \quad (5.2)$$

(ii) Let $M_Y = \{-\infty \underset{1}{\smile} 0 \underset{\circ}{\smile} 1/\lambda \underset{1}{\smile} \lambda/2 \underset{e_3(1)}{\smile} \infty\}$ (see Figure 3b). Y is a subgroup of index 3 in G_6 . An easy calculation of the side pairings shows that $Y = G_0(2) = \{(x_{ij}) \in G_6 : x_{12} \equiv 0 \pmod{2}\}$. Note that both M_X and M_Y are 3-clusters (with different side pairings).

5.2. Subgroups of index 2 of G_q . A special polygon of a subgroup of index 2 is either a 2-cluster or a union of two special triangles (1-clusters). In the case q is odd, it has to be a union of two special triangles. As a consequence, its Hecke-Farey Symbol is given as follows.

$$M_1 = \{-\infty \underset{\bullet}{\smile} 0 \underset{\bullet}{\smile} \infty\}. \quad (5.3)$$

A set of independent generators is $\{ST^{-1}, T^{-1}S\}$. In the case q is even, G_q has three subgroups of index 2, (5.3) and two more given as follows.

$$M_2 = \left\{ -\infty \underset{\circ}{\smile} 0 \underset{\circ}{\smile} 1/\lambda_q \underset{e_2(1)}{\smile} \infty \right\}, \quad M_3 = \left\{ -\infty \underset{1}{\smile} 0 \underset{1}{\smile} 1/\lambda_q \underset{e_2(1)}{\smile} \infty \right\}. \quad (5.4)$$

To determine $e_2(1)$, we once again follow the notation of (i) of subsection 4.3. One has $r = 2$, $g = 1$, $x_i = 1/\lambda_q$, and $x_{i+1} = 1/0$. It follows from (4.4) that $e_2(1) = R^2$. In particular,

$$M_3 = \left\langle R^2 = \begin{pmatrix} 0 & 1 \\ -1 & \lambda_q \end{pmatrix}^2, \begin{pmatrix} 1 & 0 \\ \lambda_q & 1 \end{pmatrix} \right\rangle. \quad (5.5)$$

M_3 is known as the *even subgroup* of G_q (q even). Note that subgroups of index 2 of G_q (q even) can be realised as Veech groups.

5.3. Commutator subgroups and principal congruence subgroups of level 2. Let $P_0 = \{x_1, x_2, \dots, x_{q-1}, x_q\}$ be the cusps of the depth 1 q -gon in the right half plane. The reduced forms of the cusps of P_0 can be determined by Lemma 3.2. Take note that $x_1 = 0$ and that $x_q = \infty$. Throughout the subsection, the x_i 's are given as above and that $x_i < x_{i+1}$.

Proposition 5.3. *A special polygon for G'_q , the commutator subgroup of G_q , is the union of two depth one q -gons forming a polygon with $2(q-1)$ free sides as follows. $G'_q \setminus \mathbb{H}$ is a genus $[(q-1)/2]$ surface with one cusp (resp. two cusps) if q is odd (resp. even).*

$$\left\{ -\infty \underset{1}{\smile} -x_{q-1} \underset{2}{\smile} \cdots \underset{q-2}{\smile} -x_2 \underset{q-1}{\smile} 0 \underset{1}{\smile} x_2 \underset{2}{\smile} \cdots \underset{q-2}{\smile} x_{q-1} \underset{q-1}{\smile} \infty \right\}. \quad (5.6)$$

Proof. The special polygon in (5.6) contains $2q$ special triangles. Hence the side pairings given by (5.6) generates a subgroup of index $2q$. As $[G_q : G'_q] = 2q$, it thus suffices to prove that the side pairings given in (5.6) are members in G'_q . This can be checked easily as the side pairing that pairs (x_i, x_{i+1}) and $(-x_{q-i+1}, -x_{q-i})$ is given by $T^{-1}R^iTR^{-i}$. $\square$

Proposition 5.4. *Let $q \geq 3$ be a prime. A special polygon for $G(2)$, the principal congruence subgroup of level 2, is the union of two depth one q -gons forming a polygon with $2(q-1)$ free sides as follows. $G(2) \setminus \mathbb{H}$ is a genus zero surface with q cusps.*

$$\left\{ -\infty \underset{1}{\smile} -x_{q-1} \underset{2}{\smile} \cdots \underset{q-2}{\smile} -x_2 \underset{q-1}{\smile} 0 \underset{q-1}{\smile} x_2 \underset{q-2}{\smile} \cdots \underset{2}{\smile} x_{q-1} \underset{1}{\smile} \infty \right\}. \quad (5.7)$$

Proof. Since q is a prime, $[G_q : G(2)] = 2q$. By (4.7), the side pairing that pairs (x_i, x_{i+1}) and $(-x_{i+1}, -x_i)$ is in $G(2)$ for every i . This completes the proof our assertion. $\square$

In the case $q \geq 3$ is a prime, similar to Proposition 5.4, one can show that $G_0(2) = G_1(2)$ has index q and admits the following special polygon (P, I_X) , where P is a q -gon.

$$\left\{ -\infty \underset{1}{\smile} x_1 \underset{2}{\smile} x_2 \cdots \underset{q-1}{\smile} x_{(q-1)/2} \underset{\circ}{\smile} x_{(q+1)/2} \cdots \underset{2}{\smile} x_{q-1} \underset{1}{\smile} \infty \right\}. \quad (5.8)$$

$G_0(2) \setminus \mathbb{H}$ is a genus zero surface with one elliptic element of order 2 and $(q+1)/2$ cusps. Note that the above is not true if q is not a prime. For instance, the group in (ii) of subsection 5.1 gives $G_0(2)$ of G_6 . Its special polygon is a 3-cluster, not a 6-gon.

5.4. Power subgroups. For each $n \in \mathbb{N}$, the power subgroup G_q^n is the characteristic subgroup of G_q generated by $\{x^n : x \in G_q\}$. Study of power subgroups has a long history back to Newman [N].

Proposition 5.5. *Let $q \geq 3$ be an odd integer. Then G_q^2 is a normal subgroup of index 2. A special polygon of G_q^2 is a union of two special triangles given as in (5.3). In the case q is even, a Hecke-Farey symbol of G_q^2 is given as in M_3 of (5.4).*

Proof. We shall first assume that q is odd. It is clear that $A = T^{-1}S$ (given as in Lemma 3.2) and $R = ST^{-1}$, each has order q , are elements of G_q^2 . Hence the special polygon of (5.3) is also a special polygon of G_q^2 .

In the case q is even, we consider the homomorphism $\phi : G_q \rightarrow \mathbb{Z}_2 = \langle a \rangle$ defined by $\phi(S) = \phi(R) = a$. It is clear that G_q^2 is the kernel of ϕ . Since G_q has exactly three subgroups of index 2 and the conjugates of S and R cannot be members of a set of independent generators of G_q^2 , we conclude that M_1 and M_2 (see (5.3) and (5.4)) are not special polygons of G_q^2 . As a consequence, a special polygon of G_q^2 is given as in M_3 of (5.4) $\square$

Proposition 5.6. *Let $\{x_1, x_2, \dots, x_{q-1}, \infty\}$ be given as in subsection 5.3 and let $r > 1$ be an odd divisor of q . Then $[G_q : G_q^r] = r$ and a Hecke-Farey symbol of G_q^r is given by*

$$\{-\infty \underset{\circ}{\smile} x_1 \underset{\circ}{\smile} x_2 \underset{\circ}{\smile} \cdots \underset{\circ}{\smile} x_r \underset{e_r(1)}{\smile} \infty\}. \quad (5.8)$$

Proof. Define $\phi : G_q \rightarrow \mathbb{Z}_r = \langle a \rangle$ by $\phi(S) = 1, \phi(R) = a$. It is clear that ϕ is a homomorphism and that $G_q^r \subseteq \ker \phi$. Hence $[G_q : G_q^r] \geq r$. Let G be the group generated by the side pairings of (5.8). The side pairings of (5.8) are conjugates of $S = S^r$ and $e_r(1) = R^r$. Hence $G \subseteq G_q^r$. Since $[G_q : G] = r$, one has $[G_q : G_q^r] \leq r$. Hence $G_q^r = G$. $\square$

5.5. Non-free normal subgroups. Let X be a proper normal subgroup of G_q that is not a free group. It follows that either $R \in X$ or $S \in X$. Suppose that $R \in X$. Since X contains all the conjugates of R , both $R = ST^{-1}$ and $T^{-1}S$ are members of X . Hence X has index 2 and a special polygon of X is given as in (5.3). Note that X is a free product of two copies of $\mathbb{Z}_q$. In the case $S \in X$, G_q/X can be generated by RX . Since R has order q , G_q/X is a cyclic group of order r , where $r|q$.

6. HURWITZ-NIELSEN REALISATION PROBLEM

6.1. Geometric invariants. An immediate application of the study of the special polygons is that the geometric invariants of $X \setminus \mathbb{H}$ can be determined easily. Let $M_X = (P, I_X)$ be a special polygon associated with $X \subseteq G_q$.

- (i) $[G_q : X]$ = the number of special triangles in $M_X = (P, I_X)$.
- (ii) The subgroup X has τ_2 (the number of the circles $\circ$ in M_X) inequivalent classes of elliptic elements of order 2 that are conjugates of S .
- (iii) The subgroup X has v_q (the number of the bullets $\bullet$ in M_X) inequivalent classes of elliptic elements of order q that are conjugates of $R = ST^{-1}$.
- (iv) Let r ($1 < r < q$) be a divisor of q . X has v_r (the number $e_{q/r}(g)$'s in M_X) inequivalent classes of elliptic elements of order r that are conjugates of $R^{q/r}$.
- (v) Suppose that the cusps of P is partitioned into v_∞ classes under the action of I_X . Then v_∞ = the number of cusps of $X \setminus \mathbb{H}$ = the number of cycles of $f(T)$, where $f(T)$ is the permutation representation of T on the set of cosets G_q/X .
- (vi) Let $\Delta = \{r : r \text{ is a divisor of } q, 2 \leq r \leq q\} = \{r_1, r_2, \dots, r_k = q\}$. The genus g of $X \setminus \mathbb{H}$ is given by the following Riemann Hurwitz formula.

$$2g - 2 + \tau_2/2 + \sum_{i=1}^k v_{r_i}(1 - 1/r_i) + v_\infty = [G_q : X](1/2 - 1/q). \quad (6.1)$$

The terms in $\{g, \tau_2, v_{r_1}, v_{r_2}, \dots, v_{r_k}, v_\infty, [G_q : X]\}$ are called the *geometric invariants* of X , where the v_i 's are given as in (ii)-(iv).

Proof. (i) is clear. (ii)-(iv) follows from the observation that each conjugacy class of elliptic elements of X must have exactly one representative in I_X as I_X is a set of independent generators. (v) follows from the fact that the number of cusps is just the number of double cosets $\langle T \rangle \setminus G_q / X$. $\square$

Lemma 6.1. *Suppose that a special polygon $M_X = (P, I_X)$ of X consists of n_0 q -gons and v_{r_i} q/r_i -clusters, $1 \leq i \leq k$. Then the number of generators of infinite order in I_X is*

$$f = \frac{n_0(q-2) + \sum_{i=1}^k v_{r_i}(q/r_i - 2) + 2 - \tau_2}{2}. \quad (6.2)$$

Proof. Since the boundary of a q/r -cluster has q/r even lines, it follows that the boundary of P has $n_0(q-2) + \sum_{i=1}^k v_{r_i}(q/r_i - 2) + 2$ even lines. Among these even lines, τ_2 of them are self paired by elements of I_X . As a consequence, the remaining even lines are free sides and they are paired by side pairings of infinite order. In particular, the number of such side pairings is $f = (n_0(q-2) + \sum_{i=1}^k v_{r_i}(q/r_i - 2) + 2 - \tau_2)/2$. $\square$

6.2. Construction of convex hyperbolic polygons. Let P_0 be either the depth one q -gon in the right half plane or the r -cluster Φ_r where $r \geq 2$ is a divisor of q (see (4.1a) for Φ_r). The Hecke-Farey sequence associated with P_0 is given by

$$F_0 = \{-\infty, 0, a_1, a_2, \dots, a_{k-1}, a_k = \infty\}. \quad (6.3)$$

Since $r \geq 2$, $\{0, a_1\}$ is an ordinary interval of F_0 . In the case P_0 is Φ_r , $\{a_{k-1}, a_k\}$ is an r -interval of F_0 (see (d) of subsection 4.3 for the terms ordinary and r -intervals). Note that P_0 is convex. Since $(0, a_1)$ is an even line of P_0 (see (d) of subsection 4.3), one may attach R_0 to P_0 along the even line $(0, a_1)$ to get a new polygon P_1 , where R_0 is either a q -gon or an s -cluster ($s \geq 2$ and $s|q$). The Hecke-Farey sequence associated with P_1 takes the form

$$F_1 = \{-\infty, 0, b_1, b_2, \dots, b_{t-1}, a_1, a_2, \dots, a_k\}. \quad (6.4)$$

Since $s \geq 2$, $\{0, b_1\}$ is an ordinary interval of P_1 and $\{b_{t-1}, a_1\}$ is an s -interval if R_0 is an s -cluster. Note that P_1 is convex. Since $(0, b_1)$ is an even line of P_1 , one may attach R_1 to P_1 along the even line $(0, b_1)$ to get a new polygon P_2 , where R_1 is either a q -gon or an u -cluster ($u \geq 2$ and $u|q$). Apply this procedure repeatedly, one admits a convex hyperbolic polygon P_n consists of n_0 q -gons and v_{r_i} q/r_i -clusters for some n_0 and v_{r_i} , where $r_i|q$ and $q/r_i \geq 2$ (the key of our construction is that P_i always has an even line $(0, x_i)$ and that R_i is always attached to P_i along $(0, x_i)$). Let $\mathbb{F}$ be the Hecke-Farey sequence associated with P_n . $\mathbb{F}$ has v_{r_i} q/r_i -intervals, where $q/r_i \geq 2$. The intervals of $\mathbb{F}$ are divided into two classes (i) q/r_i -intervals, where $q/r_i \geq 2$, and (ii) ordinary intervals. To count the number of ordinary intervals, we note that

- (i) the boundary of a q -gon has q even line,
- (ii) the boundary of a q/r_i -cluster has q/r_i even lines.

It follows that the boundary of P_n has $n_0(q-2) + \sum v_{r_i}(q/r_i - 2) + 2$ even lines. As the even lines of P_n are associated with the ordinary intervals of $\mathbb{F}$ (see (d) of subsection 4.3), the number of ordinary intervals of $\mathbb{F}$ is given by

$$n_0(q-2) + \sum v_{r_i}(q/r_i - 2) + 2, \quad q/r_i \geq 2. \quad (6.5)$$

6.3. Millington's Theorem. Kulkarni [K2, K3] gives two proofs of Millington's Theorem [Mi], one by *Diagrams* (see Section 4 of [K2]) and one by *Farey symbols* (see Section 7.6 of [K3]). We extend Millington's result to G_q by studying Hecke-Farey symbols. Our proof is a simple generalisation of Kulkarni's proof of Millington's Theorem.

Theorem 6.2. *Let $g \geq 0$, $\tau_2 \geq 0$, $v_{r_i} \geq 0$, $d \geq 1$, $v_\infty \geq 1$ be integers and let $r_i \in \Delta = \{r : r \text{ is a divisor of } q, 2 \leq r \leq q\} = \{r_1, r_2, \dots, r_k = q\}$. Then G_q has a subgroup X of index d , genus g with v_∞ cusps, and τ_2 (resp. v_{r_i}) conjugacy classes of elements that are conjugates of S (resp. R^{q/r_i}) if and only if $m_0 = 4g - 4 + \tau_2 + 2v_\infty + \sum_{i=1}^k v_{r_i}(2 - q/r_i) \geq 0$ is a multiple of $(q-2)$ and*

$$2g - 2 + \tau_2/2 + \sum_{i=1}^k v_{r_i}(1 - 1/r_i) + v_\infty = d(1/2 - 1/q). \quad (6.6)$$

Note that if q is odd, then $m_0/(q-2) \in \mathbb{Z}$ is a consequence of (6.6).

Proof. Suppose that $m_0 \geq 0$ is a multiple of $q-2$ and that $g \geq 0$, $\tau_2 \geq 0$, $v_{r_i} \geq 0$, $d \geq 1$, $v_\infty \geq 1$ satisfy the Riemann-Hurwitz formula (6.6). Let $r_i \in \Delta$. It follows that $q/r_i \geq 2$ for $1 \leq i \leq k-1$. Let $n_0 = m_0/(q-2)$.

Case 1. $n_0 = 0$ and $v_{r_i} = 0$ for all $i \leq k-1$. A simple calculation shows that either (i) $d = 1, \tau_2 = v_q = 1, v_\infty = 1, g = 0, v_{r_i} = 0$ for all $i \leq k-1$ or (ii) $d = 2, v_q = 2, v_\infty = 1, g = 0, v_{r_i} = 0$ for all $i \leq k-1$. In case (i), $X = G_q$. In case (ii), X is given as in (5.3).

Case 2. $n_0 > 0$ or $v_{r_i} > 0$ for some $i \leq k-1$. By (6.5), we have a polygon P_n and a Hecke-Farey sequence $\mathbb{F}$ with $v_{r_i} q/r_i$ -intervals ($i = 1, 2, \dots, k-1$) and

$$n_0(q-2) + \sum_{i=1}^{k-1} v_{r_i}(q/r_i - 2) + 2 = 4g - 2 + \tau_2 + v_q + 2v_\infty \geq 2 \quad (6.7)$$

ordinary intervals. Since $n_0 > 0$ or $v_{r_i} > 0$ for some $i \leq k-1$, $\mathbb{F}$ has at least three intervals. We make $\mathbb{F}$ into a Hecke-Farey symbol by declaring the first τ_2 ordinary intervals even intervals (see Definition 4.2 for the terms even and odd intervals) and the next v_q ordinary intervals odd intervals (this is equivalent to adjoin v_q special triangles to P_n). The next $2(v_\infty - 1)$ ordinary intervals are declared to be free intervals and are divided into $(v_\infty - 1)$ consecutive pairs which are paired. The remaining $4g$ ordinary intervals are free intervals and are paired in the usual $xyx^{-1}y^{-1}$ fashion (see Discussion 6.3). Let X be the group generated by the above side pairings and let I_X be the set of the side pairings.

- (i) I_X possesses τ_2 elements that are conjugates of S and v_{r_i} elements that are conjugates of R^{q/r_i} for $1 \leq i \leq k$.
- (ii) An easy study of the side pairings shows that X has v_∞ cusps. The index of X is given by the number of special triangles of the special polygon $M_X = (P, I_X)$, where P is the union of P_n and v_q special triangles. There are $n_0 = m_0/(q-2)$ q -gons, $v_{r_i} q/r_i$ -clusters ($1 \leq i \leq k-1$) and v_q special triangles. Hence the index is

$$n_0q + \sum_{i=1}^{k-1} v_{r_i}(q/r_i) + v_q = d. \quad (6.8)$$

The genus of X is g (see (6.1)). It follows that X is the required subgroup (Theorem 5.1).

Conversely, let X be given as in our theorem, the geometric invariants of X satisfy (6.6). Let $M_X = (P, I_X)$ be a special polygon of X . Then P has $v_{r_i} q/r_i$ -clusters ($1 \leq i \leq k$). Suppose that P has n_0 q -gons. By (i) of subsection 6.1, $d = n_0q + \sum_{i=1}^k v_{r_i} q/r_i$. By (6.6), one has $n_0 = (4g - 4 + \tau_2 + 2v_\infty + \sum_{i=1}^k v_{r_i}(2 - q/r_i))/(q-2)$. In particular, $4g - 4 + \tau_2 + 2v_\infty + \sum_{i=1}^k v_{r_i}(2 - q/r_i) \geq 0$ is a multiple of $q-2$. $\square$

Discussion 6.3. (i) A Hecke-Farey symbol must have at least two side pairings (subsection 4.3). (ii) Since $\mathbb{F}$ has at least three intervals, our declaration gives at least two side pairings.

6.4. Kurosh's Theorem. Let Y be a free product of F_f (a free group of rank f), π_2 copies of $\mathbb{Z}_2$, and v_{r_i} copies of $\mathbb{Z}_{r_i}$, where $r_i \in \Delta_0 = \{r : r \text{ is a divisor of } q, 3 \leq r \leq q\} = \{r_1, r_2, \dots, r_s = q\}$. We say Y is *realisable* in G_q if Y is isomorphic to a subgroup X of finite index of G_q . It is clear that $\mathbb{Z}_2 * \mathbb{Z}_q$ and $\mathbb{Z}_q * \mathbb{Z}_q$ are realisable (see (5.3) for $\mathbb{Z}_q * \mathbb{Z}_q$).

Proposition 6.4. *Let $q \geq 3$ be an odd integer and let Y be given as above. Suppose that Y is not $\mathbb{Z}_2 * \mathbb{Z}_q$ or $\mathbb{Z}_q * \mathbb{Z}_q$. Then Y is isomorphic to a subgroup X of G_q if and only if $2f + \pi_2 + v_q - 2 > 0$ and $m_0 = 2f + \pi_2 + \sum_{i=1}^s v_{r_i}(2 - q/r_i) - 2 \geq 0$ is a multiple of $q-2$. The index of X is $d = m_0q/(q-2) + \sum_{i=1}^s v_{r_i}q/r_i$.*

Proof. Suppose that Y is isomorphic to a subgroup X of G_q . Let $M_X = (P, I_X)$ be a special polygon of X . Then I_X has f elements of infinite order, π_2 elements that are conjugates of S , and v_{r_i} elements that are conjugates of R^{q/r_i} . By Lemma 6.1, $f, v_{r_i}, n_0 \geq 0$, and π_2 satisfy the identity given in (6.2). This implies that $m_0 = 2f + \pi_2 + \sum_{i=1}^s v_{r_i}(2 - q/r_i) - 2 = n_0(q-2) \geq 0$. In particular, $m_0 \geq 0$ is a multiple of $q-2$.

Note that $n_0 = m_0/(q-2)$ is the number of q -gons of P . Since P has n_0 q -gons and $v_{r_i} q/r_i$ -clusters, the index of X is $d = m_0q/(q-2) + \sum_{i=1}^s v_{r_i}q/r_i$ (see (i) of subsection 6.1). Finally, since $m_0 = 2f + \pi_2 + v_q - 2 + \sum_{i=1}^{s-1} v_{r_i}(2 - q/r_i) = n_0(q-2) \geq 0$ and $(2 - q/r_i) < 0$

for $i \leq s-1$, one has $2f + \pi_2 + v_q - 2 \geq 0$. Suppose that $2f + \pi_2 + v_q - 2 = 0$. It follows easily that $v_{r_i} = 0$ for $i \leq s-1$. A simple study shows that Y is isomorphic to a subgroup X of G_q only if $Y \cong \mathbb{Z}_2 * \mathbb{Z}_q$ or $\mathbb{Z}_q * \mathbb{Z}_q$. A contradiction. Hence $2f + \pi_2 + v_q - 2 > 0$.

Conversely, suppose that $2f + \pi_2 + v_q - 2 > 0$ and that $n_0 = m_0/(q-2) \in \mathbb{N} \cup \{0\}$. If $n_0 = 0$ and $v_{r_i} = 0$ for $i \leq s-1$, then $m_0 = 2f + \pi_2 + v_q - 2 = 0$. A contradiction. Hence either $n_0 > 0$ or $v_{r_i} > 0$ for some $i \leq s-1$. By (6.5), we have a polygon P_n and a Hecke-Farey sequence $\mathbb{F}$ with $v_{r_i} q/r_i$ -intervals ($q/r_i \geq 2$ for $i \leq s-1$) and

$$n_0(q-2) + \sum_{i=1}^{s-1} v_{r_i}(q/r_i - 2) + 2 = 2f + \pi_2 + v_q \geq 2 \quad (6.9)$$

ordinary intervals. We make $\mathbb{F}$ into a Hecke-Farey symbol by declaring the first π_2 ordinary intervals even intervals and the next v_q ordinary intervals odd intervals (this is equivalent to adjoin v_q special triangles to P_n). The last $2f$ ordinary intervals are declared to be free intervals and are divided into f consecutive pairs which are paired. Let X be the subgroup of G_q generated by the above side pairings and let I_X be the set of the side pairings. Then

- (A) I_X possesses τ_2 elements that are conjugates of S (of order 2) and v_{r_i} elements that are conjugates of R^{q/r_i} (of order r_i) for $1 \leq i \leq s$.
- (B) I_X has f elements of infinite order.

By Theorem 5.1, X is isomorphic to Y . This completes the proof of the proposition. $\square$

In the case q is even, subgroups of G_q may possess two types of elliptic elements of order 2, namely, conjugates of S and $R^{q/2}$. Note that $2 \notin \Delta_0$. For our convenience, we set $v_2 = 0$.

Proposition 6.5. *Let $q \geq 4$ be an even integer and let Y be given as above. Suppose that Y is not $\mathbb{Z}_2 * \mathbb{Z}_q$ or $\mathbb{Z}_q * \mathbb{Z}_q$. Then Y is isomorphic to a subgroup X of G_q if and only if*

- (i) $m_0 = 2f + (\pi_2 - t) + t(2 - q/2) + \sum_{i=1}^s v_{r_i}(2 - q/r_i) - 2 \geq 0$ is a multiple of $q - 2$ for some $t \geq 0$, where $(\pi_2 - t)$ is nonnegative, and
- (ii) $2f + (\pi_2 - t) + v_q - 2 \geq 0$ and $(2f + (\pi_2 - t) + v_q - 2, t, v_{q/2}) \neq (0, 0, 0)$.

Note that $v_2 = 0$. The index of X is $d = m_0 q / (q - 2) + tq/2 + \sum_{i=1}^s v_{r_i} q / r_i$.

Proof. Suppose that $Y \cong X \subseteq G_q$. Let $M_X = (P, I_X)$ be a special polygon of X . Then I_X has f elements of infinite order, $(\pi_2 - t)$ elements that are conjugates of S , t elements that are conjugates of $R^{q/2}$, and v_{r_i} elements that are conjugates of R^{q/r_i} . Note that $r_i \neq 2$ ($r_i \in \Delta_0$). The assertion now follows by applying the proof of Proposition 6.4. $\square$

7. PERMUTATION REPRESENTATION OF G_q ON G_q/X

The main purpose of this section is to give an easy and systematic method that determines the permutation representation of G_q on G_q/X .

7.1. A commutative diagram. Let Φ be given as in subsection 4.1. Φ is a fundamental domain of G_q . For each coset $Xg \in G_q/X$, there exists a unique $x_g \in X$ such that $x_g g \Phi \in M_X = (P, I_X)$. Denoted by Ω_X the set of all such special triangles $x_g g \Phi$. It follows that $\cup_{x_g g \Phi \in \Omega_X} x_g g \Phi = M_X$. As a consequence, there is a one to one correspondence between G_q/X and Ω_X defined by

$$\tau(Xg) = x_g g \Phi \in \Omega_X. \quad (7.1)$$

An element g of G_q acts on G_q/X by $(Xh, g) \rightarrow Xhg$. We shall now study the action of X on Ω_X as follows. Let $g \in G_q$. For each $g_i \Phi \in \Omega_X$, there exists a unique pair $(g_j \Phi, x_{ij}) \in \Omega_X \times X$ such that $x_{ij} g_i g \Phi = g_j \Phi \in \Omega_X$. The action of g on Ω_X is defined by

$$(g_i \Phi, g) \rightarrow x_{ij} g_i g \Phi. \quad (7.2)$$

Lemma 7.1. *The action of G_q on G_q/X is isomorphic to the action of G_q on Ω_X .*

Proof. We consider the following diagram, where the horizontal arrows represent the actions of X on G_q/X and Ω_X .

$$\begin{array}{ccc} G_q/X \times G_q & \longrightarrow & G_q/X \\ (\tau, id) \downarrow & & \downarrow \tau \\ \Omega_X \times G_q & \longrightarrow & \Omega_X \end{array} \quad (7.3)$$

Let $G_q/X = \cup Xg_i$ be chosen such that $\cup g_i\Phi = M_X$. This implies that $\tau(Xg_i) = g_i\Phi$. As a consequence, (7.3) is a commutative diagram and the action of G_q on G_q/X is isomorphic to the action of G_q on Ω_X . $\square$

7.2. The permutation representation of G_q on Ω_X . The main purpose of this subsection is to give the permutation representations of S and R on Ω_X .

Lemma 7.2. *Let $\Omega_X = \{g_i\Phi : \cup g_i\Phi = M_X\}$. The permutation representation of S on Ω_X is given by*

$$f(S) = \prod (g_i\Phi, g_j\Phi), \quad (7.4)$$

where $g_i\Phi \in \Omega_X$ and $g_j\Phi \in \Omega_X$ share the same even line of P . $(g_i\Phi, g_j\Phi)$ is a one cycle if and only if $g_iSg_i^{-1} \in X$ if and only if $g_i(0, \infty)$ is paired with itself by $g_iSg_i^{-1} \in X$.

Proof. Let $g_i\Phi \in \Omega_X$. There exists a unique pair $(g_j\Phi, x_{ij}) \in \Omega_X \times X$ such that $x_{ij}g_iS\Phi = g_j\Phi \in \Omega_X$ (see (7.2)). By Lemma 7.1, the permutation representation of S is given by $\prod(\tau(Xg_i), \tau(Xg_iS)) = \prod(g_i\Phi, x_{ij}g_iS\Phi) = \prod(g_i\Phi, g_j\Phi)$. The even line of $g_j\Phi \in M_X$ is $g_j(0, \infty) = x_{ij}g_iS(0, \infty) = x_{ij}g_i(0, \infty) \in M_X$. Since $g_i(0, \infty) \in M_X$, $x_{ij} \in X$ and every even line has a unique X -image in M_X , we conclude that $g_j(0, \infty) = x_{ij}g_i(0, \infty) = g_i(0, \infty)$. Hence $g_j\Phi$ and $g_i\Phi$ share the same even line.

$(g_i\Phi, g_j\Phi)$ is a one cycle if and only if (Xg_i, Xg_iS) is a one cycle (see Lemma 7.1 and (7.1)) if and only if $g_iSg_i^{-1} \in X$ if and only if $g_i(0, \infty)$ is paired with itself by $g_iSg_i^{-1} \in X$. $\square$

Discussion 7.3. Let $E = E_1 \cup E_2$ be the set of even lines of $M_X = (P, I_X)$, where $E_1 = \{L_1, L_2, \dots, L_m\}$ is the set of even lines that are not paired with itself by elements of I_X of order 2, and $E_2 = \{L_{m+1}, \dots, L_n\}$ is the set of even lines that are paired with itself by elements of I_X of order 2. Consequently, an element L_s of E_2 belongs to exactly one special triangle of Ω_X . Lemma 7.2 suggests a simple way to list the special triangles in Ω_X . Namely, the special triangles share the even line $L_r \in E_1$ are labeled as r and $\bar{r}$ and the special triangle contains $L_s \in E_2$ is labeled as s .

Example 7.4. Let

$$M_X = \{-\infty \underset{\bullet}{\cup} 0 \underset{\circ}{\cup} 1 \underset{1}{\cup} 2 \underset{1}{\cup} 3 \underset{\bullet}{\cup} \infty\} \quad (7.5)$$

be the Hecke-Farey symbol of a subgroup X of index 11 of $G_3 = PSL(2, \mathbb{Z})$ (see Figure 4). M_X has 6 even lines, five of them are shared by two special triangles of Ω_X , they are

$$L_1 = (0, \infty), L_2 = (1, \infty), L_3 = (2, \infty), L_4 = (3, \infty), L_5 = (1, 2). \quad (7.6)$$

Note that $(1, 2)$ and $(2, 3)$ give the same line as they are paired by the side pairing labeled by the natural number 1 (see (7.5)). The even line $L_6 = (0, 1)$ is paired to itself by an element of order 2 (see (7.5)). As a consequence, the special triangles of Ω_X are labeled as in Figure 4. By Lemma 7.2, $f(S)$ is given by

$$f(S) = (1, \bar{1})(2, \bar{2})(3, \bar{3})(4, \bar{4})(5, \bar{5})(6). \quad (7.7)$$

The permutation representation of R on G_q/X that contains $Xg_i \in \Omega_X$ is given by the cycle $(Xg_i, Xg_iR, \dots, Xg_iR^{r-1})$, where r is the smallest positive integer such that $Xg_i = Xg_iR^r$. By Lemma 7.1, the permutation representation of R on Ω_X that contains $g_i\Phi$ is given by the following r cycle.

$$(g_i\Phi = \tau(Xg_i), \tau(Xg_iR), \tau(Xg_iR^2), \dots, \tau(Xg_iR^{r-1})). \quad (7.8)$$




Figure 4

Note that $g_i(e^{\pi i/q})$ is the odd vertex of the special triangle $g_i\Phi$ and that $R(e^{\pi i/q}) = e^{\pi i/q}$. For each Xg_iR^j , by (7.2), there exists a unique pair $(g_j\Phi, x_{ij}) \in \Omega_X \times X$ such that $x_{ij}g_iR^j\Phi = g_j\Phi \in \Omega_X$. By (7.1), $\tau(Xg_iR^j) = x_{ij}g_iR^j\Phi$. The odd vertex of $x_{ij}g_iR^j\Phi$ is $x_{ij}g_iR^j(e^{\pi i/q}) = x_{ij}g_i(e^{\pi i/q}) \in M_X$. Since $x_{ij} \in X$, $g_i(e^{\pi i/q}) \in M_X$ and every odd vertex has a unique X -image in M_X , we conclude that $x_{ij}g_i(e^{\pi i/q}) = g_i(e^{\pi i/q})$. Hence the odd vertex of $\tau(Xg_iR^j) = x_{ij}g_iR^j\Phi$ is the same as the odd vertex of $g_i\Phi$ for every j . Hence the special triangles in (7.8) are special triangles of a tile Δ whose odd vertex is $g_i(e^{\pi i/q})$ (Δ is either a q -gon or an r -cluster). Since R acts as a counter-clockwise rotation about $e^{\pi i/q}$, the members in (7.8) are ordered following the orientation of M_X . The following is clear.

Lemma 7.5. *Let Δ be a tile of M_X (Δ is either a q -gon or an r -cluster). Denoted by c_Δ the cycle of special triangles of Δ ordered in the counter-clockwise manner. The permutation representation of R on Ω_X is*

$$f(R) = \prod c_\Delta. \quad (7.9)$$

Example 7.6. Let M_X be given as in (7.5). Then $f(R) = (1, 6, \bar{2})(2, 5, \bar{3})(3, \bar{5}, 4)(\bar{1})(4)$.

8. NORMALISER OF X IN $PSL(2, \mathbb{R})$

Let $G_X = \langle f(S), f(R) \rangle$. Then G_X is a subgroup of S_n , where $[G_q : X] = n$. The main purpose of this section is to determine the normaliser $N(X)$ of X in $PSL(2, \mathbb{R})$. In particular, we are able to determine whether X is a normal subgroup of G_q .

Proposition 8.1. *Let X be a subgroup of G_q of index n . Then $N_{G_q}(X)/X \cong C_{S_n}(G_X)$. X is a normal subgroup of G_q if and only if the order of G_X is $[G_q : X] = n$.*

Proof. Consider the action of G_X on Ω_X . It is clear that the action is transitive. Let G_0 be a one point stabiliser. By Lemma A1 of Appendix A, $C_{S_n}(G_X) \cong N_{G_X}(G_0)/G_0$. By Lemma 7.1, the action of G_q on G_q/X is isomorphic to the action of G_X on Ω_X . Hence $N_{G_q}(X)/X \cong N_{G_X}(G_0)/G_0 \cong C_{S_n}(G_X)$.

Consider the action of G_q on G_q/X , one has $X \triangleleft G_q$ if and only if $Y = \cap gXg^{-1} = X$ if and only if $G_X \cong G_q/Y \cong G_q/X$ if and only if G_X has order n . $\square$

Discussion 8.2. (i) Since $f(S)$ and $f(R)$ can be obtained easily by studying the special polygon M_X and the order of G_X can be determined by **GAP**, whether X is normal becomes an easy issue. In the case $q \neq 3, 4, 6$, Margulis' characterisation of arithmeticity in terms of the commensurator implies that $N(X) = N_{G_q}(X)$. Hence $C_{S_n}(G_X) \cong N(X)/X$. (ii) By Proposition 8.1, the group in (i) of subsection 5.1 is normal.

Example 8.3. Let

$$M_X = \{-\infty \underset{1}{\cup} 0 \underset{2}{\cup} 1 \underset{2}{\cup} 3/2 \underset{3}{\cup} 2 \underset{3}{\cup} 3 \underset{1}{\cup} \infty\} \quad (8.1)$$




Figure 5

be the Hecke-Farey symbol of $X \subseteq G_3 = PSL(2, \mathbb{Z})$ of index 12 (see Figure 5). M_X has 6 even lines. They are

$$L_1 = (0, \infty), L_2 = (0, 1), L_3 = (3/2, 2), L_4 = (1, \infty), L_5 = (2, \infty), L_6 = (1, 2). \quad (8.2)$$

Note that $(3, \infty)$ is paired with L_1 , $(1, 3/2)$ is paired with L_2 , and that $(2, 3)$ is paired with L_3 (see (8.1) for the side pairings). One sees easily that all the even lines are shared by 2 special triangles. Following Discussion 7.3, the special triangles are labeled as in Figure 5. By Lemmas 7.2 and 7.5, we have

$$f(R) = (1, 2, \bar{4})(4, 6, \bar{5})(5, \bar{3}, \bar{1})(\bar{6}, \bar{2}, 3), f(S) = (1, \bar{1})(2, \bar{2})(3, \bar{3})(4, \bar{4})(5, \bar{5})(6, \bar{6}). \quad (8.3)$$

One has $|G_X| = |\langle f(S), f(R) \rangle| = [G_3 : X] = 12$. By Proposition 8.1, X is a normal.

Example 8.4. Let $\lambda = \lambda_4 = \sqrt{2}$ and let

$$M_X = \{-\infty \underset{1}{\circ} 0 \underset{2}{\circ} 1 \underset{1}{\circ} \lambda \underset{3}{\circ} 3/\lambda \underset{2}{\circ} 2\lambda \underset{3}{\circ} \infty\} \quad (8.4)$$

be the Hecke-Farey symbol of $X \subseteq G_4$ of index 8 (see Figure 6). M_X has four even lines, $L_1 = (0, \infty)$, $L_2 = (0, 1/\lambda)$, $L_3 = (\lambda, 3/\lambda)$, and $L_4 = (\lambda, \infty)$. One sees from Figure 6 that

$$f(R) = (1, 2, \bar{1}, \bar{4})(4, 3, \bar{2}, \bar{3}), f(S) = (1, \bar{1})(2, \bar{2})(3, \bar{3})(4, \bar{4}). \quad (8.5)$$

One has $|G_X| = 16 > 8 = [G_4 : X]$. By Proposition 8.1, X is not normal.

Example 8.5. Let $P = \{x_1 = 0, x_2, \dots, x_q = \infty\}$ be the depth 1 q -gon in the right half plane and let r ($1 \leq r < q$) be a divisor of q . Let

$$M_A = \{-\infty \underset{\circ}{\circ} x_1 \underset{\circ}{\circ} x_2 \underset{\circ}{\circ} \dots x_{q-1} \underset{\circ}{\circ} x_q\} \text{ and } M_B = \{-\infty \underset{\circ}{\circ} x_1 \underset{\circ}{\circ} x_2 \underset{\circ}{\circ} \dots x_r \underset{\circ}{\circ} x_q\}.$$

$f(S)$ is the identity permutation for both A and B . $f(R)$ is a q -cycle (resp. r -cycle) for A (resp. B). By Proposition 8.1, both A and B are normal subgroups of G_q .

9. HURWITZ-NIELSEN REALISATION PROBLEM FOR NORMAL SUBGROUPS

A transitive subgroup G of S_d is called *regular* if every non-identity element of G is fixed point free. The result in this section reveals the connection between normal subgroups of index d of G_q and regular subgroups of S_d .

9.1. Geometric invariants revisited. Let $M_X = (P, I_X)$ be a special polygon of X . The geometric invariants of X in terms of $f(S)$ and $f(R)$ are given as follows.

- (i) I_X has τ_2 elliptic elements of order 2 that are conjugates of S , where τ_2 is the number of one cycles of $f(S)$.
- (ii) Let r ($1 < r \leq q$) be a divisor of q . I_X has v_r elliptic elements of order r that are conjugates of $R^{q/r}$, where v_r is the number of q/r -cycles of $f(R)$.
- (iii) P has v_{r_i} q/r_i -clusters. The number n_0 of q -gons of $M_X = (P, I_X)$ is the number of q -cycles of $f(R)$. The index $[G_q : X]$ is given by $n_0 q + \sum v_r q/r$.




Figure 6

(iv) The number of cusps is the number of cycles of $f(T)$, where $f(T)$ is the permutation representation of T on Ω_X (see (v) of subsection 6.1).

(v) I_X has $f = (n_0(q-2) + \sum_{r|q} v_r(q/r-2) + 2 - \tau_2)/2$ side pairings of infinite order.

Proof. By Lemma 7.5, it is clear that (a) an s -cycle ($s < q$) of $f(R)$ corresponds to an s -cluster of P as well as a side pairing which is a conjugate of R^s , (b) a q -cycle of $f(R)$ corresponds to a q -gon of P . Similarly, each one cycle of $f(S)$ corresponds to a side pairing which is a conjugate of S . (i)-(iv) of the above is clear. See Lemma 6.1 for (v). $\square$

9.2. Geometric invariants for normal subgroups. Suppose that X is normal of index d . By Proposition 8.1, $G_X = \langle f(R), f(S) \rangle$ is a transitive subgroup of order d of the symmetric group S_d . Since every nonidentity element acts freely, $f(R)$ is a product of d/u u -cycles for some $u|q$. Similarly, $f(S)$ is either 1 or a product of $d/2$ 2-cycles. The geometric invariants can be determined by subsection 9.1. In particular,

(vi) I_X has $f = d(u-2)/u + 2 - \tau_2)/2$ side pairings of infinite order ($\tau_2 = 0$ if $S \notin X$).

The following fact is well known. Unlike most of the results in the literature, our study gives the set of generators of such X 's in matrix forms (see Example 8.5).

Lemma 9.1. *Let A and B be given as in Example 8.5 and let X be a normal subgroup of index d of G_q that contains S . Then X is isomorphic to either A or B .*

Proof. Let $M_X = (P, I_X)$ and let $G_X = \langle f(S), f(R) \rangle$. Since X is normal, G_X is regular of order d . Since $S \in X$ and X is normal, $f(S) = 1$ (see (i) of subsection 9.1). Hence $G_X = \langle f(R) \rangle$. Since G_X is transitive, $f(R)$ must be a d -cycle. It follows that P is either a q -gon (if $d = q$) or a d -cluster Φ_d (if $d < q$). As $S \in X$ and X is normal, the even lines of P are self paired by conjugates of S . This completes the proof of the proposition. $\square$

9.3. Realisation of groups as normal subgroups. Throughout the subsection, Y is a free product of a free group F_f of rank f , π_2 copies of $\mathbb{Z}_2$, and v_{r_i} copies of $\mathbb{Z}_{r_i}$, where $r_i \in \Delta_0 = \{r : r|q, 3 \leq r \leq q\}$. Proposition 9.2 is an immediate consequence of Lemma 9.1.

Proposition 9.2. *Let $q \geq 3$. Suppose that $\pi_2 \neq 0$. Then Y can be realised as a normal subgroup of G_q that contains S if and only if Y is a free product of q copies of $\mathbb{Z}_2$ or a free product of $\mathbb{Z}_{q/r}$ and r copies of $\mathbb{Z}_2$, where $r|q$ and $r < q$.*

Proposition 9.3. *Let $q \geq 3$. Suppose that $\pi_2 \neq 0$. Then Y can be realised as a normal subgroup of index d of G_q that does not contain S if and only if (i) Y is a free product of $2d/q$ copies of $\mathbb{Z}_2$ and a free group of rank f , where $2f = 2d(q/2 - 2)/q + 2$, (ii) S_d has a regular subgroup $G = \langle \alpha, \beta \rangle$ of order d , where α is a product of 2-cycles and β is a product of $q/2$ -cycles. In particular, q is even. Note that d is determined by f and q .*

Proof. Let Y and G be given as in (i) and (ii). Since G_q is a free product of S and R , the map $\chi : G_q \rightarrow G$ defined by $\chi(S) = \alpha, \chi(R) = \beta$ is a homomorphism. Let X be the kernel of χ and let $M_X = (P, I_X)$ be a special polygon of X . One sees that $f(S)$ (resp. $f(R)$) and α (resp. β) have the same cycle decomposition. Hence $f(S)$ is a product of 2-cycles, $f(R)$ is

a product of $q/2$ -cycles, and P is a union of $q/2$ -gons ($2d/q$ of them). By (vi) of subsection 9.2, $2f = 2d(q/2 - 2)/q + 2$. Hence I_X has $2d/q$ elements that are conjugates of $R^{q/2}$ and f elements of infinite order. In particular, $Y \cong X \triangleleft G_q$.

Conversely, suppose that $Y \cong X \triangleleft G_q$, where $[G_q : X] = d$, $S \notin X$. Let $G_X = \langle f(S), f(R) \rangle$. By (i)-(vi) of subsections 9.1 and 9.2, G_X is regular, $f(S)$ (resp $f(R)$) is a product of 2-cycles (resp. $q/2$ -cycles). Further, X is a free product of $2d/q$ copies of $\mathbb{Z}_2$ and a free group F_f , where $2f = 2d(q/2 - 2)/q + 2$. Hence (i) and (ii) holds. $\square$

The following proposition can be proved by applying the proof of Proposition 9.3.

Proposition 9.4. *Let $q \geq 3$. Suppose that $\pi_2 = 0$. Then Y can be realised as a normal subgroup of index d of G_q if and only if*

- (i) Y is a free group of rank f , where $2f = d(q - 2)/q + 2$, S_d has regular subgroup $\langle \alpha, \beta \rangle$ of order d , where α is a product of 2-cycles and β is a product of q -cycles, or
- (ii) Y is a free product of d/r copies of $\mathbb{Z}_{q/r}$ and a free group of rank f , where $2f = d(r - 2)/r + 2$, S_d has regular subgroup $\langle \alpha, \beta \rangle$ of order d , where α is a product of 2-cycles and β is a product of r -cycles. Note that d is determined by f, q and r .

10. CONSTRUCTION OF ALL MAPS ON COMPACT ORIENTABLE SURFACES

10.1. Known results. A map $\mathfrak{M}$ on a compact orientable surface $\mathcal{X}$ is an embedding of a finite connected graph $\mathcal{G}$ in $\mathcal{X}$ such that the connected components (faces of $\mathfrak{M}$) of $\mathcal{X} \setminus \mathcal{G}$ are simply connected. An edge e of $\mathfrak{M}$ is called a *segment* if e has two vertices. An edge homeomorphic to the circle S^1 is called a *loop*. An edge homeomorphic to $[0, 1]$ with only one vertex is called a *free edge*. Segments and loops are also called *non-free edges*. The *darts* of $\mathfrak{M}$ are *directed edges*. Each non-free edge gives two darts whereas a free edge gives only one dart (see pp. 276 of [JS]). The two darts associated with a non-free edge e travel along e in opposite directions and the only dart associated with a free edge e (with vertex v) is the directed edge that points towards v . Denoted by Ω the set of darts of $\mathfrak{M}$. $\mathfrak{M}$ can be characterised completely by two permutations r_1 and r_2 on Ω . r_1 is the permutation that (i) fixes the darts associated with the free edges, and (ii) transposes the two darts associated with the non-free edges. r_2 is the permutation whose cycles correspond to the faces of $\mathfrak{M}$ (following the orientation of $\mathcal{X}$). Consequently, the map $\mathfrak{M}$ can be represented by

$$(G, \Omega, r_1, r_2), \quad (10.1)$$

where $G = \langle r_1, r_2 \rangle$. Suppose that the order of r_2 is q . Then $\theta : G_q \rightarrow G = \langle r_1, r_2 \rangle \subseteq S_\Omega$ defined by $\theta(S) = r_1$ and $\theta(R) = r_2$ is a homomorphism. Since the graph $\mathcal{G}$ is connected, θ induces a transitive action of G_q on Ω via $A(w) = \theta(A)(w)$, for all $A \in G_q$, $w \in \Omega$. Let X be a one point stabiliser of the action. It is clear that the action of G_q on G_q/X is isomorphic to the action of G on Ω and that (G, Ω, r_1, r_2) is isomorphic to $(G_q/X_0, G_q/X, S, R)$, where $X_0 = \cap_{g \in G_q} gXg^{-1}$ (see pp. 283 of [JS]). Equivalently,

$$\mathfrak{M} \cong (G, \Omega, r_1, r_2) \cong (G_q/X_0, G_q/X, S, R). \quad (10.2)$$

X is known as a *map subgroup* of $\mathfrak{M}$. Since the action is transitive, the map subgroups are conjugate to each other. See pp.63 of [CS] for more detail.

10.2. Maps associated with subgroups of G_q . Let $M_X = (P, I_X) = X \setminus \mathbb{H}$ be a special polygon of $X \subseteq G_q$ given as in Section 4. Define $M(X)$ as follows.

- (i) V = set of vertices of $M(X)$ = the set of equivalence classes of cusps of M_X ,
- (ii) E = set of edges of $M(X)$ = the set of equivalence classes of even lines of M_X ,
- (iii) F = set of faces of $M(X)$ = the set of clusters and q -gons of M_X .

As the faces of $M(X)$ are simply connected, $M(X) = (V, E, F)$ is a map.

Lemma 10.1. *Suppose that $[G_q : X] < \infty$. Then $M(X) \cong (G_X, \Omega_X, f(S), f(R))$.*

Proof. Let E_1 and E_2 be given as in Discussion 7.3. Let $e \in E_1$. Then e is not paired with itself by elements of order 2. Hence e is either a loop or a segment. Consequently, e is a non-free edge. In the case $e \in E_2$, e is paired with itself by an element of order 2. Hence e is homeomorphic to $[0, 1]$ with only one vertex. It follows that e is a free edge. Note that every non-free edge e is shared by two special triangles and every free edge e belongs to exactly one special triangle. This allows us to construct a one to one correspondence ν between the set of special triangles Ω_X and the set of darts Ω such that $\nu(f(S)(g_i\Phi)) = r_1(\nu(g_i\Phi))$ and that $\nu(f(R)(g_i\Phi)) = r_2(\nu(g_i\Phi))$. Hence the following is a commutative diagram.

$$\begin{array}{ccc} \Omega_X \times G_X & \longrightarrow & \Omega_X \\ (\nu, \chi) \downarrow & & \downarrow \nu \\ \Omega \times G & \longrightarrow & \Omega \end{array} \quad (10.3)$$

where the horizontal arrows represent the group actions and $\chi(f(S)) = r_1$, $\chi(f(R)) = r_2$. As a consequence, $M(X) \cong (G, \Omega, r_1, r_2) \cong (G_X, \Omega_X, f(S), f(R))$. $\square$

Discussion 10.2. Let $\mathfrak{M}$ be a map. One knows very little about the map subgroups of $\mathfrak{M}$ except that they are the one point stabilisers. Conversely, for each subgroup X of G_q , one cannot really *visualise* $(G_q/X_0, G_q/X, S, R)$ as a map as it is not very easy to describe the incidence relations of this map. The following proposition implies that every map (with $o(r_2) = q$) takes the form $M(X)$, where X is a subgroup of finite index of G_q . Both the map $M(X)$ and its map subgroups can be described explicitly as $M(X)$ can be described by its special polygon and the map subgroups of $M(X)$, which are conjugates of X , can be described by the set of independent generators I_X .

Proposition 10.3. *Let $\mathfrak{M}$ be a map, where $o(r_2) = q$. Then $\mathfrak{M} \cong M(X)$ for some $X \subseteq G_q$. Further, the map subgroups of $M(X)$ are conjugates of X .*

Proof. By the results in subsection 10.1, $\mathfrak{M}$ is isomorphic to $(G_q/X_0, G_q/X, S, R)$. By Lemmas 7.1, 7.2, and 7.5, $(G_q/X_0, G_q/X, S, R)$ is isomorphic to $(G_X, \Omega_X, f(S), f(R))$. By Lemma 10.1, one has $\mathfrak{M} \cong (G_X, \Omega_X, f(S), f(R)) \cong M(X)$.

$g \in G_q$ fixes $g_i\Phi \in \Omega_X$ if and only if there exists some $x \in X$ such that $xg_i\Phi = g_i\Phi$ (see (7.2)). Since the action of G_q on Ω_X is fixed point free, g fixes $g_i\Phi$ if and only if $g_i^{-1}xg_i = 1$. This completes the proof of the proposition. $\square$

Proposition 10.4. *The automorphism group of $M(X)$ is $C_{S_n}(G_X)$, where $n = [G_q : X]$.*

Proof. Let $\sigma \in S_d$. The incidence relations of $M(X)$ is determined by $G_X = \langle f(S), f(R) \rangle$. Hence $\sigma \in \text{Aut } M(X)$ if and only if $[\sigma, G_X] = 1$. This completes the proof of the lemma. $\square$

11. CONGRUENCE SUBGROUP PROBLEM FOR $\Gamma = PSL(2, \mathbb{Z})$

Let $\Gamma = G_3 = PSL(2, \mathbb{Z})$. The *principal congruence subgroup* of Γ of level $r \in \mathbb{N}$ is

$$\Gamma(r) = \{x \in \Gamma : x \equiv \pm I \pmod{r}\}. \quad (11.1)$$

$X \subseteq \Gamma$ is a *congruence subgroup* if $\Gamma(r) \subseteq X$ for some r . Consider the action of T on Γ/X , the order of T on Γ/X is called the *level* of X . By a result of Wohlfahrt [W], $X \subseteq \Gamma$ is a congruence subgroup if and only if $\Gamma(n) \subseteq X$. To the best of our knowledge, the congruence test developed by Tim Hsu [H] is the most effective one in the literature (see [LLT2] for another test). His test can be implemented as long as the permutation representations of T and U on Γ/X can be determined (Theorem 3.1 of [H]), where

$$T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, U = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}. \quad (11.2)$$

Incidentally, an algorithm that determines the permutation representations of T and U on Γ/X is not included in [H]. By Lemma 7.1, such action is isomorphic to the action of T and U on Ω_X . Following Lemmas 7.2 and 7.5, $f(T) = f(R^{-1}S)$ and $f(U) = f(RS)$




Figure 7

can be determined easily as long as X is given in terms of a special polygon and a set of independent generators. As a consequence, Hsu's algorithm can be implemented with ease.

Example 11.1. Let

$$M_X = \{-\infty \underset{\bullet}{\cup} 0 \underset{\circ}{\cup} 1 \underset{\circ}{\cup} 2 \underset{\circ}{\cup} 3 \underset{\bullet}{\cup} \infty\} \quad (11.3)$$

be the Hecke-Farey symbol of a subgroup X of index 11 of $PSL(2, \mathbb{Z})$ (see Figure 7). Then $f(S) = (1, \bar{1})(2, \bar{2})(3, \bar{3})(4, \bar{4})$ and $f(R) = (\bar{1})(1, 5, \bar{2})(2, 6, \bar{3})(3, 7, \bar{4})(4)$. Hence

$$f(T) = (\bar{1}, \bar{2}, \bar{3}, \bar{4}, 4, 7, 3, 6, 2, 5, 1), \quad f(U) = (\bar{1}, 5, \bar{2}, 6, \bar{3}, 7, \bar{4}, 4, 3, 2, 1). \quad (11.4)$$

By the algorithm given in Section 3 of [H], X is non-congruence. Note that $\cap gXg^{-1} \cong \langle f(T), f(U) \rangle \cong A_{11}$ is the alternating group on 11 letters. The group $\cap gXg^{-1}$ was first studied by Magnus [M] as part of his study of non-congruence subgroups of Γ .

APPENDIX A

In this appendix, Ω is a finite set, S_Ω is the symmetric group on Ω , and $G \subseteq S_\Omega$.

Lemma A1. *Suppose that G acts transitively on Ω . Then $C_{S_\Omega}(G) \cong N_G(G_d)/G_d$, where G_d is the one point stabiliser of $d \in \Omega$.*

Proof. Since G is transitive, The action of G on Ω is isomorphic to the action of G on the set of cosets G/G_d . Without loss of generality, we may assume that $\Omega = G/G_d$ and that $x(G_d) = xG_d$ for $x \in G$. Let $x \in C_{S_\Omega}(G)$. Then $x(G_d) = e_x G_d \in G/G_d$ for some $e_x \in G$. For each $g \in G_d$, one has $gx(G_d) = xg(G_d) = x(G_d)$. Hence $ge_x G_d = e_x G_d$. This implies that $e_x G_d \in N_G(G_d)/G_d$. As a consequence, one can show that $C_{S_\Omega}(G) \cong N_G(G_d)/G_d$ by studying the homomorphism $\Phi : C_{S_\Omega}(G) \rightarrow N_G(G_d)/G_d$ defined by $\Phi(x) = e_x^{-1} G_d$. Note that for each $r \in N_G(G_d)$, the permutation defined by $\sigma(gG_d) = gr^{-1}G_d$ commutes with G which implies that Φ is surjective ($e_\sigma = r^{-1}$ and $\Phi(\sigma) = rG_d$). $\square$

REFERENCES

- [CS] İ. N. Cangül, D. Singerman, *Normal subgroups of Hecke groups and regular maps*, Math. Proc. Camb. Phil. Soc. 123, (1998), 59-74.
- [G] GAP, *Groups, Algorithms, Programming, A system for Computational Discrete Algebra*, [http : //www.gap-system.org](http://www.gap-system.org)
- [HR] Y. H. He, J. Read, *Hecke groups, Deesins d'Enfants and the Archimedean solids*, [arXiv:math/1309.2326v1](https://arxiv.org/abs/math/1309.2326v1), [math.NT], 2013.
- [H] T. Hsu, *Identifying congruence subgroups of the modular group*, Proc. Amer. Mat. Soc. 124, no 5 (1996), 1351-1359.
- [IS] I. Ivrišimtzis, D. Singerman, *Regular maps and principal congruence subgroups of Hecke groups*, European J. of Comb. 26 (2005), 437-456.
- [JS] G. A. Jones, D. Singerman, *Theory of maps on orientable surfaces*, Proc. London Math. Soc. (3) 37 (1978), 273-307.
- [K1] R.S. Kulkarni, *An extension of a theorem of Kurosh and applications to Fuchsian groups*, Michigan Math. J. 30 (1983), 259-272.

- [K2] R.S. Kulkarni, *A new proof and extension of a theorem of Millington on the modular group*, Bull. London Math. Soc. 17 (1985), 458-462.
- [K3] R.S. Kulkarni, *An arithmetic-Geometric method in the study of subgroups of the modular group*, Amer. J Math. 113 (1991), 1053-1134.
- [LLT1] M. L. Lang, C. H. Lim, S. P. Tan, *Independent generators for congruence subgroups of Hecke groups*, Math. Z. 220 (1995), 569 – 594.
- [LLT2] M. L. Lang, C. H. Lim, S. P. Tan, *An algorithm for determining if a subgroup of the modular group is congruence*, J. of London Math. Soc. (2) 51 (1995) 491-502.
- [M] W. Magnus, *Non-Euclidean tessellations and their groups*, Academic Press 1974.
- [Ma] B. Maskit, *On Poincaré's theorem for fundamental polygons*, Advances in Math. 7, (1971), 219-230.
- [Mi] M. H. Millington, *Subgroups of the classical modular group*, J. Lon. Math. Soc. 1 (1969), 351-357.
- [N] M. Newman, *The structure of some subgroups of the Modular Group*, Illinois J. Math. (1962), 480-487.
- [W] K. Wohlfahrt, *An extension of F. Klein's level concept*, Illinois J. of Math. 8 (1964), 529 – 535.

Cheng Lien Lang

Department of Mathematics, I-Shou University, Kaohsiung, Taiwan.

`c1lang@isu.edu.tw`

Mong Lung Lang

Singapore 669608, Singapore.

`lang2to46@gmail.com`